arXiv:math/0702110v1 [math.CO] 5 Feb 2007

ON THE NUMBER OF INEQUIVALENT BINARY SELF-ORTHOGONAL CODES

XIANG-DONG HOU

ABSTRACT. Let $\Psi_{k,n}$ denote the number of inequivalent binary self-orthogonal $[n,k]$ codes. We present a method which allows us to compute $\Psi_{k,n}$ explicitly for a moderate k and an arbitrary n. Included in this paper are explicit formulas for $\Psi_{k,n}$ with $k \le 5$.

1. INTRODUCTION

Let $\mathbb{F}_2$ be the binary field. Throughout the paper, a *code* is a subspace of $\mathbb{F}_2^n$ for some integer $n > 0$; an $[n,k]$ code is a k-dimensional subspace of $\mathbb{F}_2^n$. Let $\langle \cdot, \cdot \rangle$ denote the usual inner product of $\mathbb{F}_2^n$. A code $C \subset \mathbb{F}_2^n$ is called self-orthogonal if $C \subset C^\perp$ where $C^\perp = \{x \in \mathbb{F}_2^n : \langle x, y \rangle = 0 \text{ for all } y \in C\}$. The number of $[n,k]$ self-orthogonal codes is known. Let $\Phi_{n,k}$ be the set of all $[n,k]$ self-orthogonal codes. Then

$$|\Phi_{n,k}| = \begin{cases} \dfrac{\prod_{j=1}^{k}(2^{n+1-2j}-1)}{\prod_{j=1}^{k}(2^j-1)} & \text{if } n \text{ is odd,} \\[2ex] \dfrac{(2^{n-k}-1)\prod_{j=1}^{k-1}(2^{n-2j}-1)}{\prod_{j=1}^{k}(2^j-1)} & \text{if } n \ge 2 \text{ is even,} \end{cases}$$

see MacWilliams and Sloane [9, Ch. 19, §6].

Let $\mathfrak{S}_n$ be the symmetric group on $\{1, \dots, n\}$. $\mathfrak{S}_n$ acts on $\mathbb{F}_2^n$ by permuting the coordinates of $\mathbb{F}_2^n$. Two codes $C_1, C_2 \subset \mathbb{F}_2^n$ are called equivalent if there is a $\sigma \in \mathfrak{S}_n$ such that $C_2 = C_1^\sigma$. Self-orthogonality of codes is preserved under the equivalence. Let $\Psi_{k,n}$ be the number of inequivalent $[n,k]$ self-orthogonal codes. Unlike $|\Phi_{n,k}|$, the number $\Psi_{k,n}$ is much more difficult to compute. In [1], Conway and Pless determined $\Psi_{k,2k}$ for $k \le 15$ after classifying doubly even $[32,16]$ codes. (Also see [2] for an update on $\Psi_{15,30}$.) In a recent paper [8], the author considered the asymptotic behavior of $\Psi_{k,2k}$ and proved that

$$\Psi_{k,2k} \sim \tau \cdot (2k)! 2^{\frac{1}{2}k(k-1)} \quad \text{as } k \to \infty,$$

where $\tau = \prod_{j=1}^{\infty}(1 + 2^{-j})$. However, not much else is known about the number $\Psi_{k,n}$.

The situation described above brings up a basic question: can $\Psi_{k,n}$ be computed explicitly? In this paper, we will see that the answer is "yes" for a moderate k and an arbitrary n.

Our interest in the number $\Psi_{k,n}$ is motivated by the important role it can play in the classification of $[n,k]$ self-orthogonal codes. Without knowing the number $\Psi_{k,n}$

Key words and phrases. binary self-orthogonal code, equivalence, general linear group, quadratic form, the symmetric group.

beforehand, the known method to classify $[n,k]$ self-orthogonal codes relies on the mass formula. The algorithm of this method is sketched as follows. Assume that a list of pairwise inequivalent $[n,k]$ self-orthogonal codes $C_1,\dots,C_t$ has been found and the cardinality of the automorphism group of each C_i has been determined. Then

$$n!\sum_{i=1}^{t}\frac{1}{|\mathrm{Aut}(C_i)|} \tag{1.1}$$

is the number of $[n,k]$ self-orthogonal codes equivalent to one of $C_1,\dots,C_t$. If the sum in (1.1) is $<|\Phi_{n,k}|$, search for an $[n,k]$ self-orthogonal code C_{t+1} which is inequivalent to all of $C_1,\dots,C_t$, and compute $|\mathrm{Aut}(C_{t+1})|$. Add C_{t+1} to the list $C_1,\dots,C_t$ and update the sum in (1.1). The list $C_1,\dots,C_t$ is complete when the sum in (1.1) equals $|\Phi_{n,k}|$. In fact, Conway, Pless and Sloane's classifications of self-orthogonal codes of length up to 30 and doubly even $[32,16]$ codes were obtained using this method [1, 10, 11]. On the other hand, if the number $\Psi_{k,n}$ is known beforehand, the algorithm to classify $[n,k]$ self-orthogonal codes is greatly simplified. One only has to find $\Psi_{k,n}$ pairwise inequivalent $[n,k]$ self-orthogonal codes.

A key step in the computation of $\Psi_{k,n}$ is to determine the numbers of zeros of certain quadratic forms defined on $\mathbb{F}_2^n$. This requires us to be able to tell the canonical forms of those quadratic forms. For this purpose, a brief review of canonical forms of binary quadratic forms is given in Section 2. In Section 3, we outline the method for computing $\Psi_{k,n}$. We also prove a few preliminary results to be used later. We derive the formula for $\Psi_{3,n}$ in Section 4 and the formula for $\Psi_{4,n}$ in Section 5. Most of the details of the computations in Sections 4 and 5 are included. In Section 6, we give the formula for $\Psi_{5,n}$ but omit the details of the computations. It should be clear from the paper that the method works for k beyond the range considered here. In Section 7, we give the numerical values of $\Psi_{k,n}$ for $k\le 5$ and $n\le 40$.

In our notation, $\mathbb{N}=\{0,1,2,\dots\}$. $M_{k\times n}$ is the set of all $k\times n$ matrices over $\mathbb{F}_2$. The $n\times n$ identity matrix is denoted by I_n or simply I when n is clear from the context. $\mathbf{1}(n)$ is the $1\times n$ all one vector. For two matrices A,B,

$$A\oplus B := \begin{bmatrix} A & \\ & B\end{bmatrix}.$$

For each function $f:\mathbb{F}_2^n\to\mathbb{F}_2$, $Z(f):=\{x\in\mathbb{F}_2^n : f(x)=0\}$. For $n\in\mathbb{Z}$, $\nu(n)$ is the 2-adic order of n. We also define

$$\delta(n)=\begin{cases}0 & \text{if } n\le 0,\\ 1 & \text{if } n>0.\end{cases}$$

A congruence $a\equiv b \pmod n$ is abbreviated as $a\equiv b\ (n)$.

2. Binary Quadratic Forms

A quadratic form in n variables over $\mathbb{F}_2$ is a function $f:\mathbb{F}_2^n\to\mathbb{F}_2$ of the form

$$f(x_1,\dots,x_n)=\sum_{i\le j}a_{ij}x_ix_j, \tag{2.1}$$

where $a_{ij} \in \mathbb{F}_2$. Of course, the quadratic form in (2.1) can also be written as

$$f(x_1, \dots, x_n) = \sum_{i<j} a_{ij} x_i x_j + \sum_i a_{ii} x_i.$$

Let

$$\Lambda_n = \{A \in M_{n\times n} : A^T = A \text{ and all diagonal entries of } A \text{ are } 0\}.$$

Then there is a bijection between $M_{n\times n}/\Lambda_n$ and the set of all quadratic forms in n variables over $\mathbb{F}_2$:

$$A + \Lambda_n \longleftrightarrow (x_1, \dots, x_n)A(x_1, \dots, x_n)^T.$$

We say that A is a matrix of the quadratic form $(x_1, \dots, x_n)A(x_1, \dots, x_n)^T$.

Let $f(x_1, \dots, x_n)$ and $g(x_1, \dots, x_n)$ be two quadratic forms over $\mathbb{F}_2$ with matrices A and B respectively. If $f(x_1, \dots, x_n) = g\big((x_1, \dots, x_n)Q\big)$ for some $Q \in \mathrm{GL}(n, \mathbb{F}_2)$, we say that f and g are linearly equivalent and we write $f \cong g$. Clearly, $f \cong g$ if and only if

$$A \equiv QBQ^T \pmod{\Lambda_n} \tag{2.2}$$

for some $Q \in \mathrm{GL}(n, \mathbb{F}_2)$. If two matrices $A, B \in M_{n\times n}$ satisfy (2.2), we write $A \cong B$. Therefore, finding the canonical form of a quadratic form under linear equivalence is the same as finding the canonical form of its matrix under the equivalence $\cong$. In practice, it is more convenient to work with the matrices than the quadratic forms themselves.

In the following theorem, we collect some well known results on the canonical forms and numbers of zeros of binary quadratic forms.

Theorem 2.1.

(i) *Every $A \in M_{n\times n}$ is $\cong$ equivalent to exactly one of the following canonical forms:*

$$2r\left\{\begin{bmatrix} 0&1&&&&&& \\ 0&0&&&&&& \\ &&\ddots&&&&& \\ &&&0&1&&& \\ &&&0&0&&& \\ &&&&&0&& \\ &&&&&&\ddots& \\ &&&&&&&0 \end{bmatrix}\right., \qquad 0 \le r \le \frac{n}{2},$$

$$2r\left\{\begin{bmatrix} 0&1&&&&&&&& \\ 0&0&&&&&&&& \\ &&\ddots&&&&&&& \\ &&&0&1&&&&& \\ &&&0&0&&&&& \\ &&&&&1&1&&& \\ &&&&&0&1&&& \\ &&&&&&&0&& \\ &&&&&&&&\ddots& \\ &&&&&&&&&0 \end{bmatrix}\right., \qquad 1 \le r \le \frac{n}{2},$$

$$\begin{array}{c} 2r\left\{\vphantom{\begin{matrix}0\\0\\ \ddots\\0\\0\end{matrix}}\right. \\ \vphantom{\begin{matrix}1\\0\\ \ddots\\0\end{matrix}} \end{array}\begin{bmatrix} 0&1&&&&&&&\\ 0&0&&&&&&&\\ &&\ddots&&&&&&\\ &&&0&1&&&&\\ &&&0&0&&&&\\ &&&&&1&&&\\ &&&&&&0&&\\ &&&&&&&\ddots&\\ &&&&&&&&0 \end{bmatrix}, \qquad 0\le r\le \frac{n-1}{2}.$$

We say that the matrix A and its corresponding quadratic form $f = (x_1,\dots,x_n)A(x_1,\dots,x_n)^T$ are of type $(n,r,0,0)$ or $(n,r,1,0)$ or $(n,r,0,1)$ according to the above three types of canonical forms of A. In a type (n,r,u,v), $n,r,v\in\mathbb{N}$ but $u\in\mathbb{F}_2$. The type of A or f is denoted by $\operatorname{type}(A)$ *or* $\operatorname{type}(f)$.

(ii) *We have*

$$\begin{bmatrix}1&\\&1\end{bmatrix}\cong\begin{bmatrix}1&0\\0&0\end{bmatrix},\quad \begin{bmatrix}1&1&\\0&1&\\&&1\end{bmatrix}\cong\begin{bmatrix}0&1&\\0&0&\\&&1\end{bmatrix},\quad \begin{bmatrix}1&1&&\\0&1&&\\&&1&1\\&&0&1\end{bmatrix}\cong\begin{bmatrix}0&1&&\\0&0&&\\&&0&1\\&&0&0\end{bmatrix}.$$

(iii) *If* $\operatorname{type}(A_i)=(n_i,r_i,u_i,v_i)$, $i=1,2$, *then*

$$\operatorname{type}(A_1\oplus A_2)=\bigl(n_1+n_2,\ r_1+r_2,\ u_1+u_2,\ \delta(v_1+v_2)\bigr). \tag{2.3}$$

(iv) *If* $\operatorname{type}(A)=(n,r,u,v)$ *and* $m\ge 0$, *then*

$$\operatorname{type}(A\otimes I_m)=\operatorname{type}(\underbrace{A\oplus\cdots\oplus A}_{m})=\bigl(mn,\ mr,\ mu,\ \delta(m)v\bigr). \tag{2.4}$$

(v) *Let $f(x_1,\dots,x_n)$ be a quadratic form over $\mathbb{F}_2$ of type (n,r,u,v). Then*

$$|Z(f)|=2^{n-1}+(1-v)(-1)^u2^{n-1-r}. \tag{2.5}$$

In Theorem 2.1, (i) is Dickson's theorem [3, Theorem 199] specialized for $\mathbb{F}_2$; (ii) – (v) can be verified easily; (v) is the main reason that binary quadratic forms are used in many areas, see, for example, Dillon and Dobbertin [4, Appendix A]. It is important to observe that if $\operatorname{type}(f)=(n,r,0,1)$, $|Z(f)|=2^{n-1}$ is independent of r.

Theorem 2.2. *Let $N(n,r,u,v)$ be the number of quadratic forms of type (n,r,u,v) in $x_1,\dots,x_n$ over $\mathbb{F}_2$. Then*

$$\begin{aligned} N(n,r,0,1)&=2^{r(r+1)}\frac{\prod_{i=0}^{2r}(2^{n-2r+i}-1)}{\prod_{i=1}^{r}(2^{2i}-1)},\\ N(n,r,0,0)&=2^{r^2-1}(2^r+1)\frac{\prod_{i=1}^{2r}(2^{n-2r+i}-1)}{\prod_{i=1}^{r}(2^{2i}-1)},\\ N(n,r,1,0)&=2^{r^2-1}(2^r-1)\frac{\prod_{i=1}^{2r}(2^{n-2r+i}-1)}{\prod_{i=1}^{r}(2^{2i}-1)}. \end{aligned}$$

Proof. Let $R_0(s,n)$ be the set of all polynomial functions $f(x_1,\dots,x_n)$ from $\mathbb{F}_2^n$ to $\mathbb{F}_2$ such that $\deg f\le s$ and $f(0)=0$. Let $N(n,r)$ be the number of elements in $R_0(2,n)/R_0(1,n)$ which are linearly equivalent to $x_1x_2+x_3x_4+\cdots+x_{2r-1}x_{2r}$. Then

$$N(n,r)=2^{r(r-1)}\frac{\prod_{i=1}^{2r}(2^{n-2r+i}-1)}{\prod_{i=1}^{r}(2^{2i}-1)}. \tag{2.6}$$

(Cf. [9, Ch. 15, Theorem 2] or [5, Lemma 2.2].) With a fixed r ($0\le r\le\lfloor\frac{n}{2}\rfloor$), let $N_{(n,r)}(u,v)$ be the number of $g\in R_0(1,n)$ such that

$$\text{type}(x_1x_2+x_3x_4+\cdots+x_{2r-1}x_{2r}+g)=(n,r,u,v).$$

Write $g=b_1x_1+\cdots+b_nx_n$. Then

$$\begin{aligned}&\text{type}(x_1x_2+x_3x_4+\cdots+x_{2r-1}x_{2r}+g)\\ &=\begin{cases}(n,r,0,1) & \text{if } (b_{2r+1},\dots,b_n)\ne 0,\\ (n,r,0,0) & \text{if } (b_{2r+1},\dots,b_n)=0 \text{ and } b_1b_2+b_3b_4+\cdots+b_{2r-1}b_{2r}=0,\\ (n,r,1,0) & \text{if } (b_{2r+1},\dots,b_n)=0 \text{ and } b_1b_2+b_3b_4+\cdots+b_{2r-1}b_{2r}=1.\end{cases}\end{aligned}$$

Therefore,

$$\begin{cases}N_{(n,r)}(0,1)=2^{2r}(2^{n-2r}-1),\\ N_{(n,r)}(0,0)=2^{2r-1}+2^{r-1},\\ N_{(n,r)}(1,0)=2^{2r-1}-2^{r-1}.\end{cases} \tag{2.7}$$

Since $N(n,r,u,v)=N(n,r)\cdot N_{(n,r)}(u,v)$, the conclusion of the theorem follows immediately form (2.6) and (2.7). □

We can define addition and scalar multiplication for types. Let A_1 and A_2 be square matrices over $\mathbb{F}_2$ and let $m\in\mathbb{N}$. We define

$$\text{type}(A_1)\boxplus\text{type}(A_2)=\text{type}(A_1\oplus A_2)$$

and

$$m*\text{type}(A_1)=\text{type}(\underbrace{A_1\oplus\cdots\oplus A_1}_{m})=\text{type}(A_1\otimes I_m)=\underbrace{\text{type}(A_1)\boxplus\cdots\boxplus\text{type}(A_1)}_{m}.$$

Then (2.3) and (2.4) become

$$(n_1,r_1,u_1,v_1)\boxplus(n_2,r_2,u_2,v_2)=\big(n_1+n_2,\ r_1+r_2,\ u_1+u_2,\ \delta(v_1+v_2)\big) \tag{2.8}$$

and

$$m*(n,r,u,v)=\big(mn,\ mr,\ mu,\ \delta(m)v\big). \tag{2.9}$$

In the subsequent sections, we will need to determine the number of common zeros of several quadratic forms. The following lemma is useful for this purpose.

Lemma 2.3. *Let $f_1,\dots,f_r$ be functions from $\mathbb{F}_2^n$ to $\mathbb{F}_2$. Then*

$$|Z(f_1)\cap\cdots\cap Z(f_r)|=-2^n+\frac{1}{2^{r-1}}\sum_{(a_1,\dots,a_r)\in\mathbb{F}_2^r}|Z(a_1f_1+\cdots+a_rf_r)|.$$

Proof. We have

$$\begin{aligned}
&\sum_{(a_1,\dots,a_r)\in\mathbb{F}_2^r} |Z(a_1f_1+\cdots+a_rf_r)|\\
=&\sum_{x\in\mathbb{F}_2^n}\ \sum_{\substack{(a_1,\dots,a_r)\in\mathbb{F}_2^r\\ a_1f_1(x)+\cdots+a_rf_r(x)=0}} 1\\
=&\,|Z(f_1)\cap\cdots\cap Z(f_r)|\cdot 2^r+\bigl(2^n-|Z(f_1)\cap\cdots\cap Z(f_r)|\bigr)2^{r-1}\\
=&\,|Z(f_1)\cap\cdots\cap Z(f_r)|\cdot 2^{r-1}+2^{n+r-1}.
\end{aligned}$$

It follows that

$$|Z(f_1)\cap\cdots\cap Z(f_r)|=-2^n+\frac{1}{2^{r-1}}\sum_{(a_1,\dots,a_r)\in\mathbb{F}_2^r}|Z(a_1f_1+\cdots+a_rf_r)|.$$

□

3. Outline of the Method and Preliminary Results

Let

$$S_{k\times n}=\{X\in M_{k\times n}: XX^T=0\}.$$

We will treat the elements in the symmetric group $\mathfrak{S}_n$ as $n\times n$ permutation matrices. The group $\mathrm{GL}(k,\mathbb{F}_2)\times\mathfrak{S}_n$ acts on $S_{k\times n}$ as follows: For $A\in\mathrm{GL}(k,\mathbb{F}_2)$, $P\in\mathfrak{S}_n$ and $X\in S_{k\times n}$,

$$X^{(A,P)}=A^{-1}XP.$$

Each $X\in S_{k\times n}$ generates a code (the row space of X) in $\mathbb{F}_2^n$ which is self-orthogonal and of dimension $\le k$. Two matrices $X_1,X_2\in S_{k\times n}$ generate equivalent codes if and only if X_1 and X_2 are in the same $\mathrm{GL}(k,\mathbb{F}_2)\times\mathfrak{S}_n$-orbit. Let $\Psi_{\le k,n}$ be the number of $\mathrm{GL}(k,\mathbb{F}_2)\times\mathfrak{S}_n$-orbits in $S_{k\times n}$. Then $\Psi_{\le k,n}$ is the number of inequivalent self-orthogonal codes in $\mathbb{F}_2^n$ of dimension $\le k$. Clearly,

$$\Psi_{k,n}=\Psi_{\le k,n}-\Psi_{\le k-1,n}.$$

Therefore, to compute $\Psi_{k,n}$, it suffices to compute $\Psi_{\le k,n}$. We will concentrate on $\Psi_{\le k,n}$ in the paper.

We will need two notions of equivalence between matrices. For $X_1,X_2\in M_{k\times n}$, we write $X_1\approx X_2$ if there exits $P\in\mathfrak{S}_n$ such that $X_1=X_2P$; we write $X_1\sim X_2$ if there exist $A\in\mathrm{GL}(k,\mathbb{F}_2)$ and $P\in\mathfrak{S}_n$ such that $X_1=A^{-1}X_2P$.

We first take a moment to determine $\Psi_{\le 0,n}$, $\Psi_{\le 1,n}$ and $\Psi_{\le 2,n}$. Obviously,

$$\begin{aligned}
\Psi_{\le 0,n}&=1,\\
\Psi_{\le 1,n}&=\Bigl\lfloor\frac{n}{2}\Bigr\rfloor+1.
\end{aligned}$$

Proposition 3.1. *We have*

$$\begin{aligned}
\Psi_{\le 2,n}=&\frac13\Bigl(\Bigl\lfloor\frac n6\Bigr\rfloor+1\Bigr)+\frac12\Bigl(\Bigl\lfloor\frac n2\Bigr\rfloor-\Bigl\lfloor\frac n4\Bigr\rfloor+1\Bigr)\Bigl(\Bigl\lfloor\frac n4\Bigr\rfloor+1\Bigr)\\
&+\frac1{36}\Bigl(\Bigl\lfloor\frac n2\Bigr\rfloor+3\Bigr)\Bigl(\Bigl\lfloor\frac n2\Bigr\rfloor+2\Bigr)\Bigl(\Bigl\lfloor\frac n2\Bigr\rfloor+1\Bigr).
\end{aligned}\tag{3.1}$$

Proof. Every matrix in $S_{2\times n}$ is $\sim$ equivalent to a matrix of the form

$$X_{a,b,c} := \begin{bmatrix} \overbrace{1 \cdots 1}^{a} & \overbrace{1 \cdots 1}^{b} & \overbrace{0 \cdots 0}^{c} & 0 \cdots 0 \\ 1 \cdots 1 & 0 \cdots 0 & 1 \cdots 1 & 0 \cdots 0 \end{bmatrix},$$

where a, b, c are all even. Moreover, $X_{a,b,c} \sim X_{a',b',c'}$ if and only if a', b', c' is a permutation of a, b, c. Therefore,

$$\Psi_{\le 2,n} = \bigl|\bigl\{(a_1, a_2, a_3) \in \mathbb{N}^3 : 0 \le a_1 \le a_2 \le a_3,\ a_1 + a_2 + a_3 \le \frac{n}{2}\bigr\}\bigr|.$$

Let

$$\mathcal{A} = \bigl\{(a_1, a_2, a_3) \in \mathbb{N}^3 : a_1 + a_2 + a_3 \le \frac{n}{2}\bigr\},$$
$$\mathcal{A}_{123} = \bigl\{(a, a, a) \in \mathcal{A}\bigr\}$$

and for $i, j \in \{1, 2, 3\}$, let

$$\mathcal{A}_{ij} = \{(a_1, a_2, a_3) \in \mathcal{A} : a_i = a_j\}.$$

Then

$$\begin{aligned}\Psi_{\le 2,n} &= |\mathcal{A}_{123}| + \frac{1}{3}|(\mathcal{A}_{12} \cup \mathcal{A}_{13} \cup \mathcal{A}_{23}) \setminus \mathcal{A}_{123}| + \frac{1}{6}|\mathcal{A} \setminus (\mathcal{A}_{12} \cup \mathcal{A}_{13} \cup \mathcal{A}_{23})| \\ &= \frac{2}{3}|\mathcal{A}_{123}| + \frac{1}{6}|\mathcal{A}_{12} \cup \mathcal{A}_{13} \cup \mathcal{A}_{23}| + \frac{1}{6}|\mathcal{A}|.\end{aligned}$$

By the inclusion-exclusion formula,

$$|\mathcal{A}_{12} \cup \mathcal{A}_{13} \cup \mathcal{A}_{23}| = 3|\mathcal{A}_{12}| - 2|\mathcal{A}_{123}|.$$

Hence

$$\Psi_{\le 2,n} = \frac{1}{3}|\mathcal{A}_{123}| + \frac{1}{2}|\mathcal{A}_{12}| + \frac{1}{6}|\mathcal{A}|. \tag{3.2}$$

Clearly,

$$|\mathcal{A}_{123}| = \Bigl\lfloor \frac{n}{6} \Bigr\rfloor + 1, \tag{3.3}$$

$$|\mathcal{A}| = \bigl|\bigl\{(a_1, a_2, a_3, a_4) \in \mathbb{N}^4 : a_1 + a_2 + a_3 + a_4 = \Bigl\lfloor \frac{n}{2} \Bigr\rfloor\bigr\}\bigr| = \binom{\lfloor \frac{n}{2} \rfloor + 3}{3} \tag{3.4}$$

and

$$\begin{aligned}|\mathcal{A}_{12}| &= \sum_{\substack{a_1, a_3 \in \mathbb{N} \\ 2a_1 + a_3 \le \frac{n}{2}}} 1 \\ &= \sum_{0 \le a_1 \le \lfloor \frac{n}{4} \rfloor} \ \sum_{0 \le a_3 \le \lfloor \frac{n}{2} \rfloor - 2a_1} 1 \\ &= \sum_{0 \le a_1 \le \lfloor \frac{n}{4} \rfloor} \Bigl(\Bigl\lfloor \frac{n}{2} \Bigr\rfloor - 2a_1 + 1\Bigr) \qquad (3.5) \\ &= \Bigl(\Bigl\lfloor \frac{n}{2} \Bigr\rfloor + 1\Bigr)\Bigl(\Bigl\lfloor \frac{n}{4} \Bigr\rfloor + 1\Bigr) - \Bigl\lfloor \frac{n}{4} \Bigr\rfloor \Bigl(\Bigl\lfloor \frac{n}{4} \Bigr\rfloor + 1\Bigr) \\ &= \Bigl(\Bigl\lfloor \frac{n}{2} \Bigr\rfloor - \Bigl\lfloor \frac{n}{4} \Bigr\rfloor + 1\Bigr)\Bigl(\Bigl\lfloor \frac{n}{4} \Bigr\rfloor + 1\Bigr).\end{aligned}$$

Equation (3.1) follows from (3.2) – (3.5). □

Unfortunately, the combinatorial method in Proposition 3.1 does not seem to have a generalization for $\Psi_{\le k,n}$ with $k \ge 3$. To compute $\Psi_{\le k,n}$ with $k \ge 3$, we start afresh with a more algebraic approach.

By the Burnside lemma,

$$\Psi_{\le k,n} = \frac{1}{|\mathrm{GL}(k,\mathbb{F}_2)\times\mathfrak{S}_n|}\sum_{\substack{A\in\mathrm{GL}(k,\mathbb{F}_2)\\ P\in\mathfrak{S}_n}} |\mathrm{Fix}(A,P)|, \tag{3.6}$$

where

$$\begin{aligned}\mathrm{Fix}(A,P) &= \{X\in S_{k\times n} : X^{(A,P)} = X\}\\ &= \{X\in M_{k\times n} : AX = XP,\ XX^T = 0\}.\end{aligned}$$

If G is a group, $\mathcal{C}(G)$ denotes a set of representatives of the conjugacy classes of G. For $g\in G$, $\mathrm{cent}_G(g)$ denotes the centralizer of g in G. A partition of an integer $n>0$ is a sequence of nonnegative integers $\lambda=(\lambda_1,\lambda_2,\dots)$ such that $\sum_{i\ge1} i\lambda_i = n$. We write $\lambda\vdash n$ to mean that λ is a partition of n. For $\lambda=(\lambda_1,\lambda_2,\dots)\vdash n$ and $0\le a<b$, we define

$$\lambda_{a,b} = \sum_{i\equiv a\,(b)} \lambda_i.$$

This expression will appear repreatedly in formulas later on. For each $\lambda\vdash n$, let $P_\lambda\in\mathfrak{S}_n$ be the "canonical" permutation of cycle type λ. For example, if $\lambda=(0,2,1)\vdash 7$, then $P_\lambda=(1,2)(3,4)(5,6,7)$ as a permutation and

$$P_\lambda = \begin{bmatrix}0&1\\1&0\end{bmatrix}\oplus\begin{bmatrix}0&1\\1&0\end{bmatrix}\oplus\begin{bmatrix}0&1&0\\0&0&1\\1&0&0\end{bmatrix}$$

as a permutation matrix. We can choose $\mathcal{C}(\mathfrak{S}_n)=\{P_\lambda : \lambda\vdash n\}$. If $\lambda=(\lambda_1,\lambda_2,\dots)\vdash n$, it is well known that

$$|\mathrm{cent}_{\mathfrak{S}_n}(P_\lambda)| = \lambda_1!\lambda_2!\cdots 1^{\lambda_1}2^{\lambda_2}\cdots.$$

Therefore, we can write (3.6) as

$$\begin{aligned}\Psi_{\le k,n} &= \sum_{\substack{A\in\mathcal{C}(\mathrm{GL}(k,\mathbb{F}_2))\\ P\in\mathcal{C}(\mathfrak{S}_n)}} \frac{1}{|\mathrm{cent}_{\mathrm{GL}(k,\mathbb{F}_2)}(A)|\,|\mathrm{cent}_{\mathfrak{S}_n}(P)|}|\mathrm{Fix}(A,P)|\\ &= \sum_{A\in\mathcal{C}(\mathrm{GL}(k,\mathbb{F}_2))}\frac{1}{|\mathrm{cent}_{\mathrm{GL}(k,\mathbb{F}_2)}(A)|}\sum_{\lambda=(\lambda_1,\lambda_2,\dots)\vdash n}\frac{|\mathrm{Fix}(A,P_\lambda)|}{\lambda_1!\lambda_2!\cdots1^{\lambda_1}2^{\lambda_2}\cdots}.\end{aligned} \tag{3.7}$$

In (3.7), the elements in $\mathcal{C}(\mathrm{GL}(k,\mathbb{F}_2))$ are the canonical forms of $k\times k$ invertible matrices and they can be enumerated in terms of their elementary divisors. As for $|\mathrm{cent}_{\mathrm{GL}(k,\mathbb{F}_2)}(A)|$, it suffices to assume that all the elementary divisors of A are powers of a single irreducible polynomial in $\mathbb{F}_2[x]$ since if $A=A_1\oplus A_2$, $A_i\in\mathrm{GL}(k_i,\mathbb{F}_2)$, where every elementary divisor of A_1 is prime to every elementary divisor of A_2, then

$$|\mathrm{cent}_{\mathrm{GL}(k,\mathbb{F}_2)}(A)| = |\mathrm{cent}_{\mathrm{GL}(k_1,\mathbb{F}_2)}(A_1)|\,|\mathrm{cent}_{\mathrm{GL}(k_2,\mathbb{F}_2)}(A_2)|.$$

Theorem 3.2. ([6, Theorem 3.6]) *Assume that A is a $k \times k$ matrix over $\mathbb{F}_q$ with elementary divisors $\underbrace{f^1, \dots, f^1}_{\mu_1}, \underbrace{f^2, \dots, f^2}_{\mu_2}, \dots$, where $f \in \mathbb{F}_q[x]$ is irreducible of degree d. Then*

$$|\mathrm{cent}_{\mathrm{GL}(k,\mathbb{F}_q)}(A)| = \prod_{i \geq 1} q^{d\mu_i(1\mu_1+2\mu_2+\cdots+i\mu_i+i\mu_{i+1}+\cdots)} \prod_{j=1}^{\mu_i}(1-q^{-dj}).$$

Now, the only component in (3.7) that needs to be determined is $|\mathrm{Fix}(A, P_\lambda)|$.

Theorem 3.3. *Let $\lambda = (\lambda_1, \lambda_2, \dots) \vdash n$ and let $A \in \mathrm{GL}(k, \mathbb{F}_2)$ with multiplicative order $o(A) = t$. For each $d \mid t$, let $s_d = k - \mathrm{rank}(A^d - I)$, let $B_d \in M_{k \times s_d}$ such that its columns form a basis of*

$$\{x \in \mathbb{F}_2^k : (A^d - I)x = 0\},$$

and let

$$\alpha_d = \sum_{\substack{i \geq 1,\, \nu(i) \leq \nu(t) \\ \gcd(i,t)=d}} \lambda_i. \tag{3.8}$$

Then

$$|\mathrm{Fix}(A, P_\lambda)| = 2^{\sum_{\nu(i)>\nu(t)} s_{\gcd(i,t)}\lambda_i} \cdot \mathfrak{n}(A), \tag{3.9}$$

where $\mathfrak{n}(A)$ is the number of sequences of matrices $(Y_d)_{d \mid t}$ with $Y_d \in M_{s_d \times \alpha_d}$ and

$$\sum_{d \mid t} \sum_{j=0}^{d-1} A^j B_d Y_d Y_d^T B_d^T (A^j)^T = 0. \tag{3.10}$$

Proof. First note that $|\mathrm{Fix}(A, P_\lambda)| = |\mathrm{Fix}(A, P_\lambda^{-1})|$ since P_λ and P_λ^{-1} are conjugates. We will compute $|\mathrm{Fix}(A, P_\lambda^{-1})|$ since the notation is more convenient for $|\mathrm{Fix}(A, P_\lambda^{-1})|$.

Let $X \in M_{k \times n}$. Then $AX = XP_\lambda^{-1}$ if and only if

$$X = [\dots;\ \underbrace{x_1^{(i)}, Ax_1^{(i)}, \dots, A^{i-1}x_1^{(i)}}_{i};\ \underbrace{\dots;}_{\lambda_i} \underbrace{x_{\lambda_i}^{(i)}, Ax_{\lambda_i}^{(i)}, \dots, A^{i-1}x_{\lambda_i}^{(i)}}_{i};\ \dots], \tag{3.11}$$

where $i \geq 1$ and $x_j^{(i)} \in \mathbb{F}_2^k$ with $(A^i - I)x_j^{(i)} = 0$, $1 \leq j \leq \lambda_i$. Put $d_i = \gcd(i, t)$. Note that $x_1^{(i)}, \dots, x_{\lambda_i}^{(i)}$ are in the column space of B_{d_i}, hence $[x_1^{(i)}, \dots, x_{\lambda_i}^{(i)}] = B_{d_i}X_i$ for some $X_i \in M_{s_{d_i} \times \lambda_i}$. Therefore we can write (3.11) as

$$X \approx \big[\dots;\ B_{d_i}X_i, AB_{d_i}X_i, \dots, A^{i-1}B_{d_i}X_i;\ \dots\big], \tag{3.12}$$

where $X_i \in M_{s_{d_i} \times \lambda_i}$ is arbitrary but the equivalence $\approx$ is given by a fixed column permutation. Since $(A^{d_i} - I)B_{d_i} = 0$, i.e., $A^{d_i}B_{d_i} = B_{d_i}$, we have

$$\big[\ B_{d_i}X_i, AB_{d_i}X_i, \dots, A^{i-1}B_{d_i}X_i\big] \approx \big[\ B_{d_i}X_i, AB_{d_i}X_i, \dots, A^{d_i-1}B_{d_i}X_i\big] \otimes \mathbf{1}(\frac{i}{d_i}). \tag{3.13}$$

From (3.12) and (3.13), we have

$$\begin{aligned}XX^T &= \sum_{\substack{i\ge 1\\ i/d_i \text{ odd}}} \sum_{j=0}^{d_i-1} A^j B_{d_i} X_i X_i^T B_{d_i}^T (A^j)^T\\ &= \sum_{d\mid t}\sum_{j=0}^{d-1} A^j B_d Y_d Y_d^T B_d^T (A^j)^T,\end{aligned}$$

where Y_d is the concatenation of all X_i with $\nu(i)\le\nu(t)$ and $\gcd(i,t)=d$. Thus $XX^T=0$ if and only if (3.10) holds. The equation $XX^T=0$ imposes no restriction on X_i with $\nu(i)>\nu(t)$ and the number of X_i with $\nu(i)>\nu(t)$ is

$$2^{\sum_{\nu(i)>\nu(t)} s_{d_i}\lambda_i}.$$

Therefore, we have

$$\begin{aligned}|\mathrm{Fix}(A,P_\lambda^{-1})| &= 2^{\sum_{\nu(i)>\nu(t)} s_{d_i}\lambda_i}\cdot\big(\text{the number of } (Y_d)_{d\mid t} \text{ satisfying (3.10)}\big)\\ &= 2^{\sum_{\nu(i)>\nu(t)} s_{d_i}\lambda_i}\cdot\mathfrak{n}(A).\end{aligned}$$

□

In Theorem 3.3 (equation (3.9)), the remaining question is how to compute the number $\mathfrak{n}(A)$. We now set to answer this question.

Put $Y_dY_d^T=[z_{ij}^{(d)}]_{s_d\times s_d}$ where $z_{ij}^{(d)}=z_{ji}^{(d)}$. Then each entry of the left side of (3.10) is a linear function of $z_{ij}^{(d)}$, $0\le i\le j\le s_d$, $d\mid t$. Thus (3.10) can be written as a system of L equations

$$\sum_{d\mid t}\sum_{0\le i\le j\le s_d} c_{l;ij}^{(d)} z_{ij}^{(d)}=0,\qquad 1\le l\le L, \tag{3.14}$$

where $c_{l;ij}^{(d)}\in\mathbb{F}_2$ are constants. We define $c_{l;ij}^{(d)}=0$ for $i>j$ and let $C_l^{(d)}=[c_{l;ij}^{(d)}]_{s_d\times s_d}$. Although the left side of (3.10) is a symmetric matrix of size $k\times k$, the number L of equations in (3.14) is usually much smaller than $\frac12 k(k+1)$, as we will see in actual computations.

Write $Y_d=[y_{ij}^{(d)}]_{s_d\times\alpha_d}$ and let $\widetilde{Y}_d$ be the concatenation of all the rows of Y_d, i.e.,

$$\widetilde{Y}_d=\big[y_{11}^{(d)},\dots,y_{1,\alpha_d}^{(d)},\ \dots,\ y_{s_d,1}^{(d)},\dots,y_{s_d,\alpha_d}^{(d)}\big]_{1\times(s_d\alpha_d)}.$$

Since

$$\big[y_{i1}^{(d)},\dots,y_{i,\alpha_d}^{(d)}\big]\big[y_{j1}^{(d)},\dots,y_{j,\alpha_d}^{(d)}\big]^T=z_{ij}^{(d)},$$

we have

$$\sum_{0\le i\le j\le s_d} c_{l;ij}^{(d)} z_{ij}^{(d)}=\widetilde{Y}_d(C_l^{(d)}\otimes I_{\alpha_d})\widetilde{Y}_d^T.$$

Let $Y_{1\times\theta}$ be the concatenation of all $\widetilde{Y}_d$, $d\mid t$, where

$$\theta=\sum_{d\mid t} s_d\alpha_d. \tag{3.15}$$

Then (3.14) becomes

$$Y\Big[\bigoplus_{d\mid t}(C_l^{(d)}\otimes I_{\alpha_d})\Big]Y^T=0,\qquad 1\le l\le L. \tag{3.16}$$

The left side of (3.16) is a quadratic form in Y, which is denoted by $f_l(Y)$. Therefore,

$$\begin{aligned}\mathfrak{n}(A) &= |Z(f_1)\cap\cdots\cap Z(f_L)| \\ &= -2^{\theta}+2^{-L+1}\sum_{(a_1,\dots,a_L)\in\mathbb{F}_2^L}|Z(a_1f_1+\cdots+a_Lf_L)| \qquad \text{(by Lemma 2.3).}\end{aligned}\tag{3.17}$$

In (3.17), we have

$$a_1f_1+\cdots+a_Lf_L = Y\Big[\bigoplus_{d\mid t}\Big[\Big(\sum_{l=1}^{L}a_lC_l^{(d)}\Big)\otimes I_{\alpha_d}\Big]\Big]Y^T.$$

Put

$$\mathfrak{C}_{(a_1,\dots,a_L)}=\bigoplus_{d\mid t}\Big[\Big(\sum_{l=1}^{L}a_lC_l^{(d)}\Big)\otimes I_{\alpha_d}\Big],\qquad (a_1,\dots,a_L)\in\mathbb{F}_2^L.$$

By Theorem 2.1, $|Z(a_1f_1+\cdots+a_Lf_L)|=|Z(Y\mathfrak{C}_{(a_1,\dots,a_L)}Y^T)|$ is determined by

$$\operatorname{type}(\mathfrak{C}_{(a_1,\dots,a_L)})=\boxplus_{d\mid t}\Big[\alpha_d*\operatorname{type}\Big(\sum_{l=1}^{L}a_lC_l^{(d)}\Big)\Big].$$

To sum up, we have the following algorithm for computing $\mathfrak{n}(A)$.

Algorithm 3.4 (An algorithm for computing $\mathfrak{n}(A)$)**.**

STEP 1. *Let* $Y_dY_d^T=[z_{ij}^{(d)}]$ *and write* (3.10) *entry wise, in the form of*

$$\sum_{d\mid t}\sum_{0\le i\le j\le s_d}c_{l;ij}^{(d)}z_{ij}^{(d)}=0,\qquad 1\le l\le L.$$

Record the matrices $C_l^{(d)}=[c_{l;ij}^{(d)}]$, $1\le l\le L$, $d\mid t$.

STEP 2. *For each* $(a_1,\dots,a_L)\in\mathbb{F}_2^L$ *and* $d\mid t$, *determine* $\operatorname{type}\big(\sum_{l=1}^{L}a_lC_l^{(d)}\big)$.

STEP 3. *For each* $(a_1,\dots,a_L)\in\mathbb{F}_2^L$, *put* $\mathfrak{C}_{(a_1,\dots,a_L)}=\bigoplus_{d\mid t}\big[\big(\sum_{l=1}^{L}a_lC_l^{(d)}\big)\otimes I_{\alpha_d}\big]$ *and compute*

$$\operatorname{type}(\mathfrak{C}_{(a_1,\dots,a_L)})=\boxplus_{d\mid t}\Big[\alpha_d*\operatorname{type}\Big(\sum_{l=1}^{L}a_lC_l^{(d)}\Big)\Big].$$

Find $|Z(Y\mathfrak{C}_{(a_1,\dots,a_L)}Y^T)|$ *by* (2.5).

STEP 4. *By* (3.17)

$$\mathfrak{n}(A)=-2^{\theta}+2^{-L+1}\sum_{(a_1,\dots,a_L)\in\mathbb{F}_2^L}|Z(Y\mathfrak{C}_{(a_1,\dots,a_L)}Y^T)|.\tag{3.18}$$

When k is small, the above algorithm is effective, as we will see in the following examples and in the computations of the subsequent sections.

Example 3.5. Let $A=\left[\begin{smallmatrix}0&1\\1&0\end{smallmatrix}\right]\in\mathrm{GL}(2,\mathbb{F}_2)$. We determine $\mathfrak{n}(A)$ by Algorithm 3.4. We have $o(A)=2$. Since

$$A-I=\begin{bmatrix}1&1\\1&1\end{bmatrix}\quad\text{and}\quad A^2-I=0,$$

we have $s_1 = 1$, $s_2 = 2$. We can choose

$$B_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \quad B_2 = I_2.$$

By (3.8) and (3.15),

$$\alpha_1 = \sum_{\nu(i)=0} \lambda_i = \lambda_{1,2}, \quad \alpha_2 = \sum_{\nu(i)=1} \lambda_i = \lambda_{2,4}, \quad \theta = \alpha_1 + 2\alpha_2. \tag{3.19}$$

Now we are in Step 1 of Algorithm 3.4. Put $Y_d Y_d^T = [z_{ij}^{(d)}]$. Then (3.10) becomes

$$\begin{bmatrix} z_{11}^{(1)} & z_{11}^{(1)} \\ z_{11}^{(1)} & z_{11}^{(1)} \end{bmatrix} + \begin{bmatrix} z_{11}^{(2)} + z_{22}^{(2)} & 0 \\ 0 & z_{11}^{(2)} + z_{22}^{(2)} \end{bmatrix} = 0,$$

i.e.,

$$\begin{cases} z_{11}^{(1)} = 0, \\ z_{11}^{(2)} + z_{22}^{(2)} = 0. \end{cases}$$

Thus, $L = 2$ and

$$C_1^{(1)} = [1], \qquad C_1^{(2)} = 0,$$
$$C_2^{(1)} = [0], \qquad C_2^{(2)} = I_2.$$

In Step 2 of Algorithm 3.4, we find that

$$\text{type}(a_1 C_1^{(1)} + a_2 C_2^{(1)}) = \begin{cases} (1,0,0,0) & \text{if } a_1 = 0, \\ (1,0,0,1) & \text{if } a_1 = 1, \end{cases}$$

$$\text{type}(a_1 C_1^{(2)} + a_2 C_2^{(2)}) = \begin{cases} (2,0,0,0) & \text{if } a_2 = 0, \\ (2,0,0,1) & \text{if } a_2 = 1. \end{cases}$$

In Step 3, we have

$$\begin{aligned} \text{type}(\mathfrak{C}_{(a_1,a_2)}) &= \alpha_1 * \text{type}(a_1 C_1^{(1)} + a_2 C_2^{(1)}) \boxplus \alpha_2 * \text{type}(a_1 C_1^{(2)} + a_2 C_2^{(2)}) \\ &= \begin{cases} (\theta,0,0,0) & \text{if } (a_1,a_2) = (0,0), \\ (\theta,0,0,\delta(\alpha_1)) & \text{if } (a_1,a_2) = (1,0), \\ (\theta,0,0,\delta(\alpha_2)) & \text{if } (a_1,a_2) = (0,1), \\ (\theta,0,0,\delta(\alpha_1+\alpha_2)) & \text{if } (a_1,a_2) = (1,1) \end{cases} \end{aligned}$$

and

$$|Z(Y\mathfrak{C}_{(a_1,a_2)}Y^T)| = \begin{cases} 2^\theta & \text{if } (a_1,a_2) = (0,0), \\ (2-\delta(\alpha_1))2^{\theta-1} & \text{if } (a_1,a_2) = (1,0), \\ (2-\delta(\alpha_2))2^{\theta-1} & \text{if } (a_1,a_2) = (0,1), \\ (2-\delta(\alpha_1+\alpha_2))2^{\theta-1} & \text{if } (a_1,a_2) = (1,1). \end{cases}$$

Finally, in Step 4, we arrive at

$$\begin{aligned} \mathfrak{n}(A) &= -2^\theta + 2^{-1} \cdot 2^{\theta-1}\big[8 - \delta(\alpha_1) - \delta(\alpha_2) - \delta(\alpha_1+\alpha_2)\big] \\ &= 2^{\alpha_1+2\alpha_2-2}\big[4 - \delta(\alpha_1) - \delta(\alpha_2) - \delta(\alpha_1+\alpha_2)\big]. \end{aligned}$$

By (3.9) and (3.19),

$$\begin{aligned} \text{Fix}\big(\left[\begin{smallmatrix} 0 & 1 \\ 1 & 0 \end{smallmatrix}\right], P_\lambda\big) &= 2^{2\sum_{\nu(i)>1}\lambda_i}\,\mathfrak{n}(A) \\ &= 2^{\lambda_{1,2}+2\lambda_{0,2}-2}\big[4 - \delta(\lambda_{1,2}) - \delta(\lambda_{2,4}) - \delta(\lambda_{1,2}+\lambda_{2,4})\big]. \end{aligned} \tag{3.20}$$

Example 3.6. Let $A = \left[\begin{smallmatrix} 0 & 1 \\ 1 & 1 \end{smallmatrix}\right] \in \mathrm{GL}(2, \mathbb{F}_2)$. In this cases, $o(A) = 3$ and

$$A^d - I \begin{cases} \text{is invertible} & \text{if } d = 1, \\ = 0 & \text{if } d = 3. \end{cases}$$

So, $s_1 = 0$, $s_3 = 2$, and we can choose

$$B_1 = \emptyset, \quad B_3 = I_2.$$

(Note: B_1 is a 2×0 matrix. We denote an $n \times 0$ matrix by $\emptyset$.) By (3.8),

$$\alpha_3 = \sum_{i \equiv 3\,(6)} \lambda_i = \lambda_{3,6}. \tag{3.21}$$

With $Y_d Y_d^T = [z_{ij}^{(d)}]$, (3.10) becomes

$$\begin{bmatrix} 0 & z_{11}^{(3)} + z_{12}^{(3)} + z_{22}^{(3)} \\ z_{11}^{(3)} + z_{12}^{(3)} + z_{22}^{(3)} & 0 \end{bmatrix} = 0,$$

i.e.,

$$z_{11}^{(3)} + z_{12}^{(3)} + z_{22}^{(3)} = 0.$$

Thus, $L = 1$ and

$$C_1^{(3)} = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}.$$

We have

$$\mathrm{type}(a_1 C_1^{(3)}) = \begin{cases} (2,0,0,0) & \text{if } a_1 = 0, \\ (2,1,1,0) & \text{if } a_1 = 1, \end{cases}$$

$$\mathrm{type}(\mathfrak{C}_{a_1}) = \alpha_3 * \mathrm{type}(a_1 C_1^{(3)}) = \begin{cases} (2\alpha_3, 0, 0, 0) & \text{if } a_1 = 0, \\ (2\alpha_3, \alpha_3, \alpha_3, 0) & \text{if } a_1 = 1 \end{cases}$$

and

$$|Z(Y \mathfrak{C}_{a_1} Y^T)| = \begin{cases} 2^{2\alpha_3} & \text{if } a_1 = 0, \\ 2^{2\alpha_3 - 1} + (-1)^{\alpha_3} 2^{\alpha_3 - 1} & \text{if } a_1 = 1. \end{cases}$$

Therefore,

$$\mathfrak{n}(A) = -2^{2\alpha_3} + 2^{2\alpha_3} + 2^{2\alpha_3 - 1} + (-1)^{\alpha_3} 2^{\alpha_3 - 1} = 2^{2\alpha_3 - 1} + (-1)^{\alpha_3} 2^{\alpha_3 - 1}.$$

By (3.9) and (3.21),

$$\begin{aligned} \mathrm{Fix}\left(\left[\begin{smallmatrix} 0 & 1 \\ 1 & 1 \end{smallmatrix}\right], P_\lambda\right) &= 2^{2 \sum_{i \equiv 0\,(6)} \lambda_i}\, \mathfrak{n}(A) \\ &= 2^{2\lambda_{0,6} - 1} \big[2^{2\lambda_{3,6}} + (-1)^{\lambda_{3,6}} 2^{\lambda_{3,6}} \big]. \end{aligned} \tag{3.22}$$

For the rest of this section, we collect a few lemmas which will simplify the computation of $|\mathrm{Fix}(A, P_\lambda)|$ in many cases.

Lemma 3.7. *We have*

$$|\mathrm{Fix}(I_k, P_\lambda)| = 2^{k\lambda_{0,2}} \Big| S_{k \times \lambda_{1,2}} \Big|. \tag{3.23}$$

Proof. Clearly, $o(I_k) = 1$, $s_1 = k$ and we can choose $B_1 = I_k$. By (3.8), $\alpha_1 = \sum_{\nu(i)=0} \lambda_i = \lambda_{1,2}$. Equation (3.10) becomes

$$Y_1 Y_1^T = 0,$$

where $Y_1 \in M_{k\times\alpha_1}$. The number of such Y_1, i.e. $\mathfrak{n}(I_k)$, is $|S_{k\times\alpha_1}|$. Hence by (3.9),

$$|\mathrm{Fix}(I_k, P_\lambda)| = 2^{k\sum_{\nu(i)>0}\lambda_i}\big|S_{k\times\alpha_1}\big|.$$

$\square$

In (3.23), $\big|S_{k\times\lambda_{1,2}}\big|$ is given by the next lemma.

Lemma 3.8. *We have*

$$|S_{k\times n}| = \sum_{0\le l\le \min\{k,\frac{n}{2}\}} |\Phi_{n,l}|\, 2^{\frac12 l(l-1)} \prod_{j=k-l+1}^{k} (2^j - 1). \tag{3.24}$$

Proof. For each $0 \le l \le \min\{k, \frac{n}{2}\}$, let

$$S_{k\times n}^{(l)} = \{X \in S_{k\times n} : \operatorname{rank} X = l\}.$$

Put

$$\mathcal{X} = \{(X, C) : X \in S_{k\times n}^{(l)},\ C \in \Phi_{n,l},\ C \text{ is the row space of } X\}.$$

Counting the number of elements $(X, C) \in \mathcal{X}$ in the order of X, C and in the order of C, X, we have

$$|S_{k\times n}^{(l)}| = |\Phi_{n,l}|\,|\{A \in M_{k\times l} : \operatorname{rank} A = l\}|.$$

In fact, for each $X \in S_{k\times n}^{(l)}$, there is a unique $C \in \Phi_{n,l}$ such that $(X, C) \in \mathcal{X}$. On the other hand, for each $C \in \Phi_{n,l}$ with a basis $c_1, \dots, c_l$, $(X, C) \in \mathcal{X}$ if and only if

$$X = A \begin{bmatrix} c_1 \\ \vdots \\ c_l \end{bmatrix}$$

for some $A \in M_{k\times l}$ with $\operatorname{rank} A = l$. Thus

$$\begin{aligned} |S_{k\times n}^{(l)}| &= |\Phi_{n,l}|(2^k - 2^0)(2^k - 2^1)\cdots(2^k - 2^{l-1}) \\ &= |\Phi_{n,l}|\, 2^{\frac12 l(l-1)} \prod_{j=k-l+1}^{k} (2^j - 1). \end{aligned}$$

It follows that

$$\begin{aligned} |S_{k\times n}| &= \sum_{0\le l\le \min\{k,\frac{n}{2}\}} |S_{k\times n}^{(l)}| \\ &= \sum_{0\le l\le \min\{k,\frac{n}{2}\}} |\Phi_{n,l}|\, 2^{\frac12 l(l-1)} \prod_{j=k-l+1}^{k} (2^j - 1). \end{aligned}$$

$\square$

We take another look of Lemmas 3.7 and 3.8. Let $\alpha_1 = \lambda_{1,2}$. Then $|S_{k\times\alpha_1}| = \mathfrak{n}(I_k)$ and Lemma 3.8 gives a formula for this number in terms of $|\Phi_{n,l}|$. However, $\mathfrak{n}(I_k)$ can also be computed directly using Algorithm 3.4, resulting in a formula for $\mathfrak{n}(I_k)$ not involving $|\Phi_{n,l}|$. We follow the notation of Algorithm 3.4. With $Y_1Y_1^T = [z_{ij}^{(1)}]_{k\times k}$, (3.10) becomes

$$z_{ij}^{(1)} = 0, \qquad 1 \le i \le j \le k.$$

So, $L = \frac{1}{2}k(k+1)$ and

$$C_1^{(1)} = \begin{bmatrix} 1 & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \end{bmatrix},\ C_2^{(1)} = \begin{bmatrix} 0 & 1 & \cdots & 0 \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \end{bmatrix},\ \dots,\ C_{k+1}^{(1)} = \begin{bmatrix} 0 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \end{bmatrix},$$

$$\dots,\ C_{\frac{1}{2}k(k+1)}^{(1)} = \begin{bmatrix} 0 & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{bmatrix}.$$

As $(a_1, \dots, a_{\frac{1}{2}k(k+1)})$ runs through $\mathbb{F}_2^{\frac{1}{2}k(k+1)}$,

$$\text{type}(a_1C_1^{(1)} + \cdots + a_{\frac{1}{2}k(k+1)}C_{\frac{1}{2}k(k+1)}^{(1)}) = (k, r, u, v) \qquad N(k, r, u, v) \text{ times},$$

where $N(k, r, u, v)$ is given by Theorem 2.2. Thus,

$$\text{type}\big(\mathfrak{C}_{(a_1,\dots,a_{\frac{1}{2}k(k+1)})}\big) = \big(k\alpha_1,\ r\alpha_1,\ u\alpha_1,\ \delta(\alpha_1)v\big) \qquad N(k, r, u, v) \text{ times}.$$

Therefore,

$$\begin{aligned}
&|S_{k\times\alpha_1}| \\
=&\mathfrak{n}(I_k) \\
=&-2^{k\alpha_1} + 2^{-\frac{1}{2}k(k+1)+1}\Big[\sum_{0\le r\le\frac{k}{2}} N(k, r, 0, 0)(2^{k\alpha_1-1} + 2^{k\alpha_1-1-r\alpha_1}) \\
&\quad + \sum_{1\le r\le\frac{k}{2}} N(k, r, 1, 0)\big(2^{k\alpha_1-1} + (-1)^{\alpha_1}2^{k\alpha_1-1-r\alpha_1}\big) \\
&\quad + \sum_{0\le r\le\frac{k-1}{2}} N(k, r, 0, 1)\big(2^{k\alpha_1-1} + (1-\delta(\alpha_1))2^{k\alpha_1-1}\big)\Big] \\
=&-2^{k\alpha_1} + 2^{-\frac{1}{2}(k+2)(k-1)} \sum_{0\le r\le\frac{k}{2}} \frac{\prod_{i=1}^{2r}(2^{k-2r+i}-1)}{\prod_{i=1}^{r}(2^{2i}-1)}\, 2^{r^2-1} \\
&\quad \cdot \Big[2^{k\alpha_1+r} + 2^{k\alpha_1-1-r\alpha_1}\big(2^r + 1 + (-1)^{\alpha_1}(2^r-1)\big) \\
&\quad + 2^{r+1}(2^{k-2r}-1)\big(2^{k\alpha_1} - \delta(\alpha_1)2^{k\alpha_1-1}\big)\Big].
\end{aligned} \tag{3.25}$$

Lemma 3.9. *Let $P \in \mathfrak{S}_n$, $A \in \text{GL}(k, \mathbb{F}_2)$ and $B \in \text{GL}(l, \mathbb{F}_2)$. Assume that the characteristic polynomials of A and B^{-1} are relatively prime. Then*

$$\text{Fix}(A \oplus B, P) = \Big\{\begin{bmatrix} X \\ Y \end{bmatrix} : X \in \text{Fix}(A, P),\ Y \in \text{Fix}(B, P)\Big\}. \tag{3.26}$$

In particular,

$$|\mathrm{Fix}(A\oplus B,P)|=|\mathrm{Fix}(A,P)|\,|\mathrm{Fix}(B,P)|. \tag{3.27}$$

Proof. It suffices to prove that in (3.26), the right hand side is contained in the left hand side. Let $X\in\mathrm{Fix}(A,P)$ and $Y\in\mathrm{Fix}(B,P)$. We have $XX^T=0$, $YY^T=0$, and

$$\begin{bmatrix}A&\\&B\end{bmatrix}\begin{bmatrix}X\\Y\end{bmatrix}=\begin{bmatrix}X\\Y\end{bmatrix}P. \tag{3.28}$$

It follows that

$$XY^T=(XP)(YP)^T=(AX)(BY)^T=AXY^TB^T,$$

i.e.,

$$A(XY^T)=(XY^T)(B^{-1})^T.$$

Since the characteristic polynomials of A and $(B^{-1})^T$ are relatively prime, we have $XY^T=0$. Thus $\left[\begin{smallmatrix}X\\Y\end{smallmatrix}\right]\in S_{(k+l)\times n}$, which, combined with (3.28), implies that $\left[\begin{smallmatrix}X\\Y\end{smallmatrix}\right]\in\mathrm{Fix}(A\oplus B,P)$. □

Corollary 3.10. *Let $f_1,\dots,f_t\in\mathbb{F}_2[x]\setminus\{x\}$ be irreducible such that $\{f_1,f_1^*\},\dots,\{f_t,f_t^*\}$ are pairwise disjoint, where f_i^* is the reciprocal polynomial of f_i. Let $A=A_1\oplus\cdots\oplus A_t\in\mathrm{GL}(k,\mathbb{F}_2)$, where $A_i\in\mathrm{GL}(k_i,\mathbb{F}_2)$ whose elementary divisors are powers of f_i or f_i^*. Then for each $P\in\mathfrak{S}_n$,*

$$|\mathrm{Fix}(A,P)|=\prod_{i=1}^t|\mathrm{Fix}(A_i,P)|.$$

Lemma 3.11. *Let $f\in\mathbb{F}_2[x]\setminus\{x\}$ be an irreducible polynomial which is not self-reciprocal. Let t be the smallest positive integer such that $f\mid x^t-1$. Let $A\in\mathrm{GL}(k,\mathbb{F}_2)$ have elementary divisors $\underbrace{f^1,\dots,f^1}_{\mu_1},\dots,\underbrace{f^s,\dots,f^s}_{\mu_s}$ and let $\lambda=(\lambda_1,\lambda_2,\dots)\vdash n$. Then*

$$|\mathrm{Fix}(A,P_\lambda)|=|\{X\in M_{k\times n}:AX=XP_\lambda\}|=2^{\deg f\sum_{j\ge1}\lambda_{jt}\sum_{l\ge1}\mu_l\min\{l,2^{\nu(j)}\}}. \tag{3.29}$$

Proof. Since the characteristic polynomials of A and A^{-1} are relatively prime, by the proof of Lemma 3.9, $AX=XP_\lambda$ implies $XX^T=0$. Hence

$$\mathrm{Fix}(A,P_\lambda)=\{X\in M_{k\times n}:AX=XP_\lambda\}.$$

To see the second equality in (3.29), note from (3.11) that

$$\dim\{X\in M_{k\times n}:AX=XP_\lambda\}=\sum_{i\ge1}\bigl[k-\mathrm{rank}\,(A^i-I)\bigr]\lambda_i.$$

By [7, Lemma 5.2 and its proof], we have

$$k-\mathrm{rank}\,(A^i-I)=\begin{cases}\deg f\displaystyle\sum_{l\ge1}\mu_l\min\{l,2^{\nu(i)}\} & \text{if } t\mid i,\\ 0 & \text{if } t\nmid i.\end{cases}$$

Therefore,

$$\dim\{X\in M_{k\times n}:AX=XP_\lambda\}=\deg f\sum_{j\ge1}\lambda_{jt}\sum_{l\ge1}\mu_l\min\{l,2^{\nu(j)}\}.$$

Table 1. Information about $\mathcal{C}(\mathrm{GL}(3,\mathbb{F}_2))$

representative	elementary divisors	$\lvert\mathrm{cent}_{\mathrm{GL}(3,\mathbb{F}_2)}(\ )\rvert$
$A_1 = I_3$	$x+1,\ x+1,\ x+1$	$2^3\cdot 3\cdot 7$
$A_2 = [1]\oplus\begin{bmatrix}0&1\\1&0\end{bmatrix}$	$x+1,\ (x+1)^2$	2^3
$A_3 = \begin{bmatrix}0&1&0\\0&0&1\\1&1&1\end{bmatrix}$	$(x+1)^3$	2^2
$A_4 = [1]\oplus\begin{bmatrix}0&1\\1&1\end{bmatrix}$	$x+1,\ x^2+x+1$	3
$A_5 = \begin{bmatrix}0&1&0\\0&0&1\\1&1&0\end{bmatrix}$	x^3+x+1	7
$A_6 = \begin{bmatrix}0&1&0\\0&0&1\\1&0&1\end{bmatrix}$	x^3+x^2+1	7

□

4. Computation of $\Psi_{\le 3,n}$

Recall that $\mathcal{C}(\mathrm{GL}(k,\mathbb{F}_2))$ is a set of representatives of the conjugacy classes of $\mathrm{GL}(k,\mathbb{F}_2)$. We let $\mathcal{C}(\mathrm{GL}(3,\mathbb{F}_2)) = \{A_1,\dots,A_6\}$ where $A_1,\dots,A_6$ are given in Table 1. By (3.7), to determine $\Psi_{\le 3,n}$, it suffices to determine $|\mathrm{Fix}(A_i,P_\lambda)|$ for $i=1,\dots,6$ and for all $\lambda=(\lambda_1,\lambda_2,\dots)\vdash n$. In the following computations, we will use the notation of Theorem 3.3 and Algorithm 3.4.

4.1. Computation of $|\mathrm{Fix}(A_1,P_\lambda)|$.

By Lemma 3.7 and (3.25),

$$\begin{aligned}|\mathrm{Fix}(A_1,P_\lambda)| &= 2^{3\lambda_{0,2}}\left|S_{3\times\lambda_{1,2}}\right|\\ &= 2^{3\lambda_{0,2}}\left[2^{3\lambda_{1,2}-6}\big(36-35\delta(\lambda_{1,2})\big)+2^{2\lambda_{1,2}-6}\,7\big(3+(-1)^{\lambda_{1,2}}\big)\right].\end{aligned}$$

4.2. Computation of $|\mathrm{Fix}(A_2,P_\lambda)|$.

We have $o(A_2)=2$. Since

$$A_2 - I = [0]\oplus\begin{bmatrix}1&1\\1&1\end{bmatrix},\qquad\text{and}\qquad A_2^2 - I = 0,$$

we have $s_1=2$, $s_2=3$, and we can choose

$$B_1 = [1]\oplus\begin{bmatrix}1\\1\end{bmatrix},\quad B_2 = I_3.$$

By (3.8) and (3.15),

$$\alpha_1=\sum_{\nu(i)=0}\lambda_i=\lambda_{1,2},\quad \alpha_2=\sum_{\nu(i)=1}\lambda_i=\lambda_{2,4},\quad \theta=2\alpha_1+3\alpha_2. \tag{4.1}$$

With $Y_dY_d^T = [z_{ij}^{(d)}]$, equation (3.10) becomes

$$\begin{bmatrix} z_{11}^{(1)} & z_{12}^{(1)} & z_{12}^{(1)} \\ z_{12}^{(1)} & z_{22}^{(1)} & z_{22}^{(1)} \\ z_{12}^{(1)} & z_{22}^{(1)} & z_{22}^{(1)} \end{bmatrix} + \begin{bmatrix} 0 & z_{12}^{(2)} + z_{13}^{(2)} & z_{12}^{(2)} + z_{13}^{(2)} \\ z_{12}^{(2)} + z_{13}^{(2)} & z_{22}^{(2)} + z_{33}^{(2)} & 0 \\ z_{12}^{(2)} + z_{13}^{(2)} & 0 & z_{22}^{(2)} + z_{33}^{(2)} \end{bmatrix} = 0,$$

i.e.,

$$\begin{cases} z_{11}^{(1)} = z_{22}^{(1)} = 0, \\ z_{12}^{(1)} + z_{12}^{(2)} + z_{13}^{(2)} = 0, \\ z_{22}^{(2)} + z_{33}^{(2)} = 0. \end{cases}$$

Thus, $L = 4$ and

$$C_1^{(1)} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \qquad C_1^{(2)} = 0,$$
$$C_2^{(1)} = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}, \qquad C_2^{(2)} = 0,$$
$$C_3^{(1)} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \qquad C_3^{(2)} = \begin{bmatrix} 0 & 1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix},$$
$$C_4^{(1)} = 0, \qquad C_4^{(2)} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

We have

$$\text{type}(a_1C_1^{(1)} + \cdots + a_4C_4^{(1)}) = \begin{cases} (2,0,0,0) & \text{if } (a_1,a_2,a_3) = (0,0,0), \\ (2,0,0,1) & \text{if } a_3 = 0,\ (a_1,a_2) \neq (0,0), \\ (2,1,0,0) & \text{if } a_3 = 1,\ (a_1,a_2) \neq (1,1), \\ (2,1,1,0) & \text{if } (a_1,a_2,a_3) = (1,1,1), \end{cases}$$

$$\text{type}(a_1C_1^{(2)} + \cdots + a_4C_4^{(2)}) = \begin{cases} (3,0,0,0) & \text{if } (a_3,a_4) = (0,0), \\ (3,0,0,1) & \text{if } (a_3,a_4) = (0,1), \\ (3,1,0,0) & \text{if } a_3 = 1. \end{cases}$$

As $(a_1,\dots,a_4)$ runs over $\mathbb{F}_2^4$,

$$\begin{aligned} &\text{type}(\mathfrak{C}_{(a_1,\dots,a_4)}) \\ &= \alpha_1 * \text{type}(a_1C_1^{(1)} + \cdots + a_4C_4^{(1)}) \boxplus \alpha_2 * \text{type}(a_1C_1^{(2)} + \cdots + a_4C_4^{(2)}) \\ &= \begin{cases} (\theta,0,0,0) & 1 \text{ time}, \\ (\theta,0,0,\delta(\alpha_2)) & 1 \text{ time}, \\ (\theta,0,0,\delta(\alpha_1)) & 3 \text{ times}, \\ (\theta,0,0,\delta(\alpha_1+\alpha_2)) & 3 \text{ times}, \\ (\theta,\alpha_1+\alpha_2,0,0) & 6 \text{ times}, \\ (\theta,\alpha_1+\alpha_2,\alpha_1,0) & 2 \text{ times}. \end{cases} \end{aligned}$$

Therefore,

$$\begin{aligned}\mathfrak{n}(A_2) = &-2^{\theta} + 2^{-3}\Big[2^{\theta} + 2^{\theta-1} + (1-\delta(\alpha_2))2^{\theta-1} + 3\big[2^{\theta-1} + (1-\delta(\alpha_1))2^{\theta-1}\big]\\ &+ 3\big[2^{\theta-1} + (1-\delta(\alpha_1+\alpha_2))2^{\theta-1}\big] + 6\big(2^{\theta-1} + 2^{\theta-1-(\alpha_1+\alpha_2)}\big)\\ &+ 2\big[2^{\theta-1} + (-1)^{\alpha_1}2^{\theta-1-(\alpha_1+\alpha_2)}\big]\Big]\\ =&\,2^{\theta-4}\big[8 - 3\delta(\alpha_1) - \delta(\alpha_2) - 3\delta(\alpha_1+\alpha_2)\big] + 2^{\alpha_1+2\alpha_2-3}(3+(-1)^{\alpha_1}).\end{aligned}$$

By (3.9) and (4.1),

$$\begin{aligned}&|\mathrm{Fix}(A_2, P_\lambda)|\\ =&\,2^{3\sum_{\nu(i)>1}\lambda_i}\,\mathfrak{n}(A_2)\\ =&\,2^{3\lambda_{0,4}}\Big[2^{2\lambda_{1,2}+3\lambda_{2,4}-4}\big[8 - 3\delta(\lambda_{1,2}) - \delta(\lambda_{2,4}) - 3\delta(\lambda_{1,2}+\lambda_{2,4})\big]\\ &+ 2^{\lambda_{1,2}+2\lambda_{2,4}-3}(3+(-1)^{\lambda_{1,2}})\Big].\end{aligned}$$

4.3. **Computation of** $|\mathrm{Fix}(A_3, P_\lambda)|$**.**

We have $o(A_3) = 4$. Since

$$A_3 - I = \begin{bmatrix}1&1&0\\0&1&1\\1&1&0\end{bmatrix}, \qquad A_3^2 - I = \begin{bmatrix}1&0&1\\1&0&1\\1&0&1\end{bmatrix}, \qquad A_3^4 - I = 0,$$

we have $s_1 = 1, s_2 = 2, s_4 = 3$ and we can choose

$$B_1 = \begin{bmatrix}1\\1\\1\end{bmatrix}, \quad B_2 = \begin{bmatrix}1&0\\0&1\\1&0\end{bmatrix}, \quad B_4 = I_3.$$

By (3.8) and (3.15)

$$\alpha_1 = \lambda_{1,2}, \quad \alpha_2 = \lambda_{2,4}, \quad \alpha_4 = \lambda_{4,8}, \quad \theta = \alpha_1 + 2\alpha_2 + 3\alpha_4. \tag{4.2}$$

With $Y_dY_d^T = [z_{ij}^{(d)}]$, equation (3.10) becomes

$$\begin{bmatrix}z_{11}^{(1)}&z_{11}^{(1)}&z_{11}^{(1)}\\z_{11}^{(1)}&z_{11}^{(1)}&z_{11}^{(1)}\\z_{11}^{(1)}&z_{11}^{(1)}&z_{11}^{(1)}\end{bmatrix} + \begin{bmatrix}z_{11}^{(2)}+z_{22}^{(2)}&0&z_{11}^{(2)}+z_{22}^{(2)}\\0&z_{11}^{(2)}+z_{22}^{(2)}&0\\z_{11}^{(2)}+z_{22}^{(2)}&0&z_{11}^{(2)}+z_{22}^{(2)}\end{bmatrix}$$

$$+\begin{bmatrix}0&z_{11}^{(4)}+z_{33}^{(4)}&0\\z_{11}^{(4)}+z_{33}^{(4)}&0&z_{11}^{(4)}+z_{33}^{(4)}\\0&z_{11}^{(4)}+z_{33}^{(4)}&0\end{bmatrix} = 0,$$

i.e.,

$$\begin{cases}z_{11}^{(1)} + z_{11}^{(2)} + z_{22}^{(2)} = 0,\\ z_{11}^{(1)} + z_{11}^{(4)} + z_{33}^{(4)} = 0.\end{cases}$$

Thus, $L = 2$ and

$$C_1^{(1)} = [1], \qquad C_1^{(2)} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \qquad C_1^{(4)} = 0,$$

$$C_2^{(1)} = [1], \qquad C_2^{(2)} = 0, \qquad C_2^{(4)} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

We have

$$\text{type}(a_1 C_1^{(1)} + a_2 C_2^{(1)}) = \begin{cases} (1,0,0,0) & \text{if } a_1 = a_2, \\ (1,0,0,1) & \text{if } a_1 \neq a_2, \end{cases}$$

$$\text{type}(a_1 C_1^{(2)} + a_2 C_2^{(2)}) = \begin{cases} (2,0,0,0) & \text{if } a_1 = 0, \\ (2,0,0,1) & \text{if } a_1 = 1, \end{cases}$$

$$\text{type}(a_1 C_1^{(4)} + a_2 C_2^{(4)}) = \begin{cases} (3,0,0,0) & \text{if } a_2 = 0, \\ (3,0,0,1) & \text{if } a_2 = 1, \end{cases}$$

$$\text{type}(\mathfrak{C}_{(a_1,a_2)}) = \begin{cases} (\theta,0,0,0) & \text{if } (a_1,a_2) = (0,0), \\ (\theta,0,0,\delta(\alpha_2+\alpha_4)) & \text{if } (a_1,a_2) = (1,1), \\ (\theta,0,0,\delta(\alpha_1+\alpha_2)) & \text{if } (a_1,a_2) = (1,0), \\ (\theta,0,0,\delta(\alpha_1+\alpha_4)) & \text{if } (a_1,a_2) = (0,1). \end{cases}$$

Thus,

$$\begin{aligned} \mathfrak{n}(A_3) = & -2^\theta + 2^{-1}\Big[2^\theta + 2^{\theta-1} + \big(1-\delta(\alpha_2+\alpha_4)\big)2^{\theta-1} \\ & + 2^{\theta-1} + \big(1-\delta(\alpha_1+\alpha_2)\big)2^{\theta-1} + 2^{\theta-1} + \big(1-\delta(\alpha_1+\alpha_4)\big)2^{\theta-1}\Big] \\ = & \, 2^{\theta-2}\big[4 - \delta(\alpha_1+\alpha_2) - \delta(\alpha_1+\alpha_4) - \delta(\alpha_2+\alpha_4)\big]. \end{aligned}$$

By (3.9) and (4.2),

$$\begin{aligned} & |\text{Fix}(A_3, P_\lambda)| \\ = & \, 2^{3\sum_{\nu(i)>2}\lambda_i}\,\mathfrak{n}(A_3) \\ = & \, 2^{\lambda_{1,2}+2\lambda_{2,4}+3\lambda_{0,4}-2}\big[4 - \delta(\lambda_{1,2}+\lambda_{2,4}) - \delta(\lambda_{1,2}+\lambda_{4,8}) - \delta(\lambda_{2,4}+\lambda_{4,8})\big]. \end{aligned}$$

4.4. **Computation of $|\text{Fix}(A_4, P_\lambda)|$.**

By Corollary 3.10,

$$|\text{Fix}(A_4, P_\lambda)| = |\text{Fix}([1], P_\lambda)|\,\big|\text{Fix}(\left[\begin{smallmatrix} 0 & 1 \\ 1 & 1 \end{smallmatrix}\right], P_\lambda)\big|,$$

where $\big|\text{Fix}(\left[\begin{smallmatrix} 0 & 1 \\ 1 & 1 \end{smallmatrix}\right], P_\lambda)\big|$ is given by (3.22). By Lemma 3.7,

$$|\text{Fix}([1], P_\lambda)| = 2^{\lambda_{0,2}}\big|S_{1\times\lambda_{1,2}}\big| = 2^{\lambda_{0,1}-1}\big(2-\delta(\lambda_{1,2})\big). \tag{4.3}$$

Therefore,

$$|\text{Fix}(A_4, P_\lambda)| = 2^{\lambda_{0,1}+2\lambda_{0,6}-2}\big(2-\delta(\lambda_{1,2})\big)\Big[2^{2\lambda_{3,6}} + (-1)^{\lambda_{3,6}}2^{\lambda_{3,6}}\Big].$$

4.5. Computation of $|\mathrm{Fix}(A_5, P_\lambda)|$ and $|\mathrm{Fix}(A_6, P_\lambda)|$.

The smallest positive integer t such that $x^3+x+1 \mid x^t-1$ is 7. By Lemma 3.11,

$$|\mathrm{Fix}(A_5, P_\lambda)| = 2^{3\lambda_{0,7}}. \tag{4.4}$$

In the same way,

$$|\mathrm{Fix}(A_6, P_\lambda)| = 2^{3\lambda_{0,7}}.$$

5. Computation of $\Psi_{\le 4,n}$

We let $\mathcal{C}\big(\mathrm{GL}(4,\mathbb{F}_2)\big) = \{A_1,\dots,A_{14}\}$ where $A_1,\dots,A_{14}$ are given in Table 2. It suffices to determine $|\mathrm{Fix}(A_i, P_\lambda)|$ for $i=1,\dots,14$ and for all $\lambda=(\lambda_1,\lambda_2,\dots)\vdash n$. Again, in the following computations, we will use the notation of Theorem 3.3 and Algorithm 3.4.

5.1. Computation of $|\mathrm{Fix}(A_1, P_\lambda)|$.

By Lemma 3.7 and (3.25),

$$\begin{aligned}
|\mathrm{Fix}(A_1, P_\lambda)| &= 2^{4\lambda_{0,2}}\big|S_{4\times\lambda_{1,2}}\big| \\
&= 2^{4\lambda_{0,2}}\Big[2^{4\lambda_{1,2}-10}\big(436-435\delta(\lambda_{1,2})\big) \\
&\quad + 2^{3\lambda_{1,2}-10}\,35\big(3+(-1)^{\lambda_{1,2}}\big) + 2^{2\lambda_{1,2}-7}\,7\big(5+3(-1)^{\lambda_{1,2}}\big)\Big].
\end{aligned} \tag{5.1}$$

5.2. Computation of $|\mathrm{Fix}(A_2, P_\lambda)|$.

We have $o(A_2)=2$ and

$$A_2 - I = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \oplus \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}, \quad A_2^2 - I = 0.$$

Hence $s_1 = 3,\ s_2 = 4$. We can choose

$$B_1 = I_2 \oplus \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \quad B_2 = I_4.$$

By (3.8) and (3.15),

$$\alpha_1 = \lambda_{1,2}, \quad \alpha_2 = \lambda_{2,4}, \quad \theta = 3\alpha_1 + 4\alpha_2. \tag{5.2}$$

With $Y_dY_d^T = [z_{ij}^{(d)}]$, equation (3.10) becomes

$$\begin{bmatrix} z_{11}^{(1)} & z_{12}^{(1)} & z_{13}^{(1)} & z_{13}^{(1)} \\ z_{12}^{(1)} & z_{22}^{(1)} & z_{23}^{(1)} & z_{23}^{(1)} \\ z_{13}^{(1)} & z_{23}^{(1)} & z_{33}^{(1)} & z_{33}^{(1)} \\ z_{13}^{(1)} & z_{23}^{(1)} & z_{33}^{(1)} & z_{33}^{(1)} \end{bmatrix} + \begin{bmatrix} 0 & 0 & z_{13}^{(2)}+z_{14}^{(2)} & z_{13}^{(2)}+z_{14}^{(2)} \\ 0 & 0 & z_{23}^{(2)}+z_{24}^{(2)} & z_{23}^{(2)}+z_{24}^{(2)} \\ z_{13}^{(2)}+z_{14}^{(2)} & z_{23}^{(2)}+z_{24}^{(2)} & z_{33}^{(2)}+z_{44}^{(2)} & 0 \\ z_{13}^{(2)}+z_{14}^{(2)} & z_{23}^{(2)}+z_{24}^{(2)} & 0 & z_{33}^{(2)}+z_{44}^{(2)} \end{bmatrix} = 0,$$

i.e.,

$$\begin{cases} z_{11}^{(1)} = z_{22}^{(1)} = z_{33}^{(1)} = z_{12}^{(1)} = 0, \\ z_{13}^{(1)} + z_{13}^{(2)} + z_{14}^{(2)} = 0, \\ z_{23}^{(1)} + z_{23}^{(2)} + z_{24}^{(2)} = 0, \\ z_{33}^{(2)} + z_{44}^{(2)} = 0. \end{cases}$$

Table 2. Information about $\mathcal{C}(\mathrm{GL}(4,\mathbb{F}_2))$

representative	elementary divisors	$\lvert\mathrm{cent}_{\mathrm{GL}(4,\mathbb{F}_2)}(\ )\rvert$
$A_1 = I_4$	$x+1,\ x+1,\ x+1,\ x+1$	$2^6\cdot 3^2\cdot 5\cdot 7$
$A_2 = I_2 \oplus \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$	$x+1,\ x+1,\ (x+1)^2$	$2^6\cdot 3$
$A_3 = [1] \oplus \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 1 & 1 \end{bmatrix}$	$x+1,\ (x+1)^3$	2^4
$A_4 = I_2 \oplus \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix}$	$x+1,\ x+1,\ x^2+x+1$	$2\cdot 3^2$
$A_5 = [1] \oplus \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix}$	$x+1,\ x^3+x+1$	7
$A_6 = [1] \oplus \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 1 \end{bmatrix}$	$x+1,\ x^3+x^2+1$	7
$A_7 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \oplus \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$	$(x+1)^2,\ (x+1)^2$	$2^5\cdot 3$
$A_8 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \oplus \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix}$	$(x+1)^2,\ x^2+x+1$	$2\cdot 3$
$A_9 = \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix} \oplus \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix}$	$x^2+x+1,\ x^2+x+1$	$2^2\cdot 3^2\cdot 5$
$A_{10} = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \end{bmatrix}$	$(x+1)^4$	2^3
$A_{11} = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 \end{bmatrix}$	$(x^2+x+1)^2$	$2^2\cdot 3$
$A_{12} = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix}$	$x^4+x^3+x^2+x+1$	$3\cdot 5$
$A_{13} = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 1 & 0 & 0 \end{bmatrix}$	x^4+x+1	$3\cdot 5$
$A_{14} = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 \end{bmatrix}$	x^4+x^3+1	$3\cdot 5$

Thus, $L = 7$ and

$$C_1^{(1)} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \qquad C_1^{(2)} = 0,$$

$$C_2^{(1)} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \qquad C_2^{(2)} = 0,$$

$$C_3^{(1)} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \qquad C_3^{(2)} = 0,$$

$$C_4^{(1)} = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \qquad C_4^{(2)} = 0,$$

$$C_5^{(1)} = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \qquad C_5^{(2)} = \begin{bmatrix} 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix},$$

$$C_6^{(1)} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}, \qquad C_6^{(2)} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix},$$

$$C_7^{(1)} = 0, \qquad C_7^{(2)} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

Fix $(a_4, a_5, a_6, a_7) \in \mathbb{F}_2^4$ and let (a_1, a_2, a_3) run over $\mathbb{F}_2^3$. If $(a_4, a_5, a_6) = (0, 0, 0)$,

$$\operatorname{type}(a_1 C_1^{(1)} + \cdots + a_7 C_7^{(1)}) = \begin{cases} (3, 0, 0, 0) & 1 \text{ time}, \\ (3, 0, 0, 1) & 7 \text{ times}. \end{cases}$$

If $(a_4, a_5, a_6) \neq (0, 0, 0)$,

$$\operatorname{type}(a_1 C_1^{(1)} + \cdots + a_7 C_7^{(1)}) = \begin{cases} (3, 1, 0, 0) & 3 \text{ times}, \\ (3, 1, 1, 0) & 1 \text{ time}, \\ (3, \square, 0, 1) & 4 \text{ times}. \end{cases}$$

In the above, the symbol $\square$ represents a component of a type which is not needed in the subsequent computations. We also have

$$\operatorname{type}(a_1 C_1^{(2)} + \cdots + a_7 C_7^{(2)}) = \begin{cases} (4, 0, 0, 0) & \text{if } (a_5, a_6, a_7) = (0, 0, 0), \\ (4, 0, 0, 1) & \text{if } (a_5, a_6, a_7) = (0, 0, 1), \\ (4, 1, 0, 0) & \text{if } (a_5, a_6) \neq (0, 0). \end{cases}$$

Therefore, as $(a_1,\dots,a_7)$ runs over $\mathbb{F}_2^7$,

$$
\begin{aligned}
&\text{type}(\mathfrak{C}_{(a_1,\dots,a_7)})\\
=&\,\alpha_1 * \text{type}(a_1C_1^{(1)}+\cdots+a_7C_7^{(1)}) \boxplus \alpha_2 * \text{type}(a_1C_1^{(2)}+\cdots+a_7C_7^{(2)})\\
=&\begin{cases}
(\theta,0,0,0) & 1 \text{ time},\\
(\theta,0,0,\delta(\alpha_2)) & 1 \text{ time},\\
(\theta,\alpha_1\square,0,\delta(\alpha_1)) & 11 \text{ times},\\
(\theta,\alpha_1\square,0,\delta(\alpha_1+\alpha_2)) & 11 \text{ times},\\
(\theta,\alpha_1,0,0) & 3 \text{ times},\\
(\theta,\alpha_1,0,\delta(\alpha_2)) & 3 \text{ times},\\
(\theta,\alpha_1+\alpha_2,0,0) & 2^2\cdot 3^2 \text{ times},\\
(\theta,\alpha_1,\alpha_1,0) & 1 \text{ time},\\
(\theta,\alpha_1,\alpha_1,\delta(\alpha_2)) & 1 \text{ time},\\
(\theta,\alpha_1+\alpha_2,\alpha_1,0) & 2^2\cdot 3 \text{ times},\\
(\theta,\alpha_1\square+\alpha_2,0,\delta(\alpha_1)) & 2^4\cdot 3 \text{ times}.
\end{cases}
\end{aligned}
$$

By (3.18),

$$
\begin{aligned}
\mathfrak{n}(A_2) = {}& 2^{3\alpha_1+4\alpha_2-7}\big[24-\delta(\alpha_2)-11\delta(\alpha_1)-11\delta(\alpha_1+\alpha_2)\big]\\
&+2^{2\alpha_1+4\alpha_2-7}\big(3+(-1)^{\alpha_1}\big)\big(2-\delta(\alpha_2)\big)\\
&+2^{2\alpha_1+3\alpha_2-5}\big[7+(-1)^{\alpha_1}-4\delta(\alpha_1)\big].
\end{aligned}
$$

By (3.9) and (5.2),

$$
\begin{aligned}
&|\text{Fix}(A_2,P_\lambda)|\\
=&\,2^{4\sum_{\nu(i)>1}\lambda_i}\,\mathfrak{n}(A_2)\\
=&\,2^{4\lambda_{0,4}}\Big[2^{3\lambda_{1,2}+4\lambda_{2,4}-7}\big[24-\delta(\lambda_{2,4})-11\delta(\lambda_{1,2})-11\delta(\lambda_{1,2}+\lambda_{2,4})\big]\\
&+2^{2\lambda_{1,2}+4\lambda_{2,4}-7}\big(3+(-1)^{\lambda_{1,2}}\big)\big(2-\delta(\lambda_{2,4})\big)\\
&+2^{2\lambda_{1,2}+3\lambda_{2,4}-5}\,3\big[7+(-1)^{\lambda_{1,2}}-4\delta(\lambda_{1,2})\big]\Big].
\end{aligned}
$$

5.3. **Computation of $|\text{Fix}(A_3,P_\lambda)|$.**

The data for the matrix A_3 are as follows: $o(A_3)=4$, $s_1=2$, $s_2=3$, $s_4=4$,

$$
B_1=[1]\oplus\begin{bmatrix}1\\1\\1\end{bmatrix},\quad B_2=[1]\oplus\begin{bmatrix}1&0\\0&1\\1&0\end{bmatrix},\quad B_4=I_4,
$$

$$
\alpha_1=\lambda_{1,2},\quad \alpha_2=\lambda_{2,4},\quad \alpha_4=\lambda_{4,8},\quad \theta=2\alpha_1+3\alpha_2+4\alpha_4. \tag{5.3}
$$

With $Y_d Y_d^T = [z_{ij}^{(d)}]$, equation (3.10) becomes

$$\begin{bmatrix} z_{11}^{(1)} & z_{12}^{(1)} & z_{12}^{(1)} & z_{12}^{(1)} \\ z_{12}^{(1)} & z_{22}^{(1)} & z_{22}^{(1)} & z_{22}^{(1)} \\ z_{12}^{(1)} & z_{22}^{(1)} & z_{22}^{(1)} & z_{22}^{(1)} \\ z_{12}^{(1)} & z_{22}^{(1)} & z_{22}^{(1)} & z_{22}^{(1)} \end{bmatrix} + \begin{bmatrix} 0 & z_{12}^{(2)} + z_{13}^{(2)} & z_{12}^{(2)} + z_{13}^{(2)} & z_{12}^{(2)} + z_{13}^{(2)} \\ z_{12}^{(2)} + z_{13}^{(2)} & z_{22}^{(2)} + z_{33}^{(2)} & 0 & z_{22}^{(2)} + z_{33}^{(2)} \\ z_{12}^{(2)} + z_{13}^{(2)} & 0 & z_{22}^{(2)} + z_{33}^{(2)} & 0 \\ z_{12}^{(2)} + z_{13}^{(2)} & z_{22}^{(2)} + z_{33}^{(2)} & 0 & z_{22}^{(2)} + z_{33}^{(2)} \end{bmatrix}$$

$$+ \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & z_{22}^{(4)} + z_{44}^{(4)} & 0 \\ 0 & z_{22}^{(4)} + z_{44}^{(4)} & 0 & z_{22}^{(4)} + z_{44}^{(4)} \\ 0 & 0 & z_{22}^{(4)} + z_{44}^{(4)} & 0 \end{bmatrix} = 0,$$

i.e.,

$$\begin{cases} z_{11}^{(1)} = 0, \\ z_{12}^{(1)} + z_{12}^{(2)} + z_{13}^{(2)} = 0, \\ z_{22}^{(1)} + z_{22}^{(2)} + z_{33}^{(2)} = 0, \\ z_{22}^{(1)} + z_{22}^{(4)} + z_{44}^{(4)} = 0. \end{cases}$$

Thus, $L = 4$ and

$$C_1^{(1)} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \qquad C_1^{(2)} = 0, \qquad C_1^{(4)} = 0,$$

$$C_2^{(1)} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \qquad C_2^{(2)} = \begin{bmatrix} 0 & 1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \qquad C_2^{(4)} = 0,$$

$$C_3^{(1)} = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}, \qquad C_3^{(2)} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \qquad C_3^{(4)} = 0,$$

$$C_4^{(1)} = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}, \qquad C_4^{(2)} = 0, \qquad C_3^{(4)} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

We have

$$\begin{aligned} &\text{type}(a_1 C_1^{(1)} + \cdots + a_4 C_4^{(1)}) \\ &= \begin{cases} (2,0,0,0) & \text{if } a_2 = 0,\ (a_1, a_3, a_4) = (0,0,0), (0,1,1), \\ (2,0,0,1) & \text{if } a_2 = 0,\ (a_1, a_3, a_4) \neq (0,0,0), (0,1,1), \\ (2,1,1,0) & \text{if } a_2 = 1,\ (a_1, a_3, a_4) = (1,0,1), (1,1,0), \\ (2,1,0,0) & \text{if } a_2 = 0,\ (a_1, a_3, a_4) \neq (1,0,1), (1,1,0), \end{cases} \end{aligned}$$

$$\text{type}(a_1 C_1^{(2)} + \cdots + a_4 C_4^{(2)}) = \begin{cases} (3,0,0,0) & \text{if } (a_2, a_3) = (0,0), \\ (3,0,0,1) & \text{if } (a_2, a_3) = (0,1), \\ (3,1,0,0) & \text{if } a_2 = 1, \end{cases}$$

$$\text{type}(a_1 C_1^{(4)} + \cdots + a_4 C_4^{(4)}) = \begin{cases} (4,0,0,0) & \text{if } a_4 = 0, \\ (4,0,0,1) & \text{if } a_4 = 1. \end{cases}$$

As $(a_1,\dots,a_4)$ runs over $\mathbb{F}_2^4$,

$$\operatorname{type}(\mathfrak{C}_{(a_1,\dots,a_4)})=\begin{cases}(\theta,0,0,0) & 1\text{ time},\\ (\theta,0,0,\delta(\alpha_2+\alpha_4)) & 1\text{ time},\\ (\theta,0,0,\delta(\alpha_1)) & 1\text{ time},\\ (\theta,0,0,\delta(\alpha_1+\alpha_4)) & 2\text{ times},\\ (\theta,0,0,\delta(\alpha_1+\alpha_2)) & 2\text{ times},\\ (\theta,0,0,\delta(\alpha_1+\alpha_2+\alpha_4)) & 1\text{ time},\\ (\theta,\alpha_1+\alpha_2,\alpha_1,0) & 1\text{ time},\\ (\theta,\alpha_1+\alpha_2,\alpha_1,\delta(\alpha_4)) & 1\text{ time},\\ (\theta,\alpha_1+\alpha_2,0,0) & 3\text{ times},\\ (\theta,\alpha_1+\alpha_2,0,\delta(\alpha_4)) & 3\text{ times}.\end{cases}$$

By (3.18),

$$\begin{aligned}\mathfrak{n}(A_3)=\,&2^{2\alpha_1+3\alpha_2+4\alpha_4-4}\big[8-\delta(\alpha_2+\alpha_4)-\delta(\alpha_1)-2\delta(\alpha_1+\alpha_4)-2\delta(\alpha_1+\alpha_2)\\ &-\delta(\alpha_1+\alpha_2+\alpha_4)\big]+2^{\alpha_1+2\alpha_2+4\alpha_4-4}\big(3+(-1)^{\alpha_1}\big)\big(2-\delta(\alpha_4)\big).\end{aligned}$$

By (3.9) and (5.3),

$$\begin{aligned}&|\mathrm{Fix}(A_3,P_\lambda)|\\ =\,&2^{4\sum_{\nu(i)>2}\lambda_i}\,\mathfrak{n}(A_3)\\ =\,&2^{4\lambda_{0,8}}\Big[2^{2\lambda_{1,2}+3\lambda_{2,4}+4\lambda_{4,8}-4}\big[8-\delta(\lambda_{2,4}+\lambda_{4,8})-\delta(\lambda_{1,2})\\ &-2\delta(\lambda_{1,2}+\lambda_{4,8})-2\delta(\lambda_{1,2}+\lambda_{2,4})-\delta(\lambda_{1,2}+\lambda_{2,4}+\lambda_{4,8})\big]\\ &+2^{\lambda_{1,2}+2\lambda_{2,4}+4\lambda_{4,8}-4}\big(3+(-1)^{\lambda_{1,2}}\big)\big(2-\delta(\lambda_{4,8})\big)\Big].\end{aligned}$$

5.4. **Computation of $|\mathrm{Fix}(A_4,P_\lambda)|$.**

By Corollary 3.10, (3.22), (3.23) and (3.25), we have

$$\begin{aligned}|\mathrm{Fix}(A_4,P_\lambda)|&=|\mathrm{Fix}(I_2,P_\lambda)|\,\big|\mathrm{Fix}\big(\left[\begin{smallmatrix}0&1\\1&1\end{smallmatrix}\right],P_\lambda\big)\big|\\ &=2^{2\lambda_{0,2}+2\lambda_{0,6}-1}\big(2^{2\lambda_{3,6}}+(-1)^{\lambda_{3,6}}2^{\lambda_{3,6}}\big)\\ &\quad\cdot\big[2^{2\lambda_{1,2}-3}\big(4-3\delta(\lambda_{1,2})\big)+2^{\lambda_{1,2}-3}\big(3+(-1)^{\lambda_{1,2}}\big)\big].\end{aligned}$$

5.5. **Computation of $|\mathrm{Fix}(A_5,P_\lambda)|$ and $|\mathrm{Fix}(A_6,P_\lambda)|$.**

Write $A_5=[1]\oplus C$ where

$$C=\begin{bmatrix}0&1&0\\0&0&1\\1&1&0\end{bmatrix}.$$

By Corollary 3.10,

$$|\mathrm{Fix}(A_5,P_\lambda)|=|\mathrm{Fix}([1],P_\lambda)|\,|\mathrm{Fix}(C,P_\lambda)|,$$

where $|\mathrm{Fix}([1],P_\lambda)|$ is given by (4.3) and $|\mathrm{Fix}(C,P_\lambda)|$ is given by (4.4). Hence we have

$$|\mathrm{Fix}(A_5,P_\lambda)|=2^{\lambda_{0,1}+3\lambda_{0,7}-1}\big(2-\delta(\lambda_{1,2})\big).$$

In the same way,

$$|\mathrm{Fix}(A_6,P_\lambda)|=2^{\lambda_{0,1}+3\lambda_{0,7}-1}\big(2-\delta(\lambda_{1,2})\big).$$

5.6. **Computation of $|\mathrm{Fix}(A_7, P_\lambda)|$.**

The data for A_7 are as follows: $o(A_7) = 2$, $s_1 = 2$, $s_2 = 4$,

$$B_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \oplus \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \quad B_2 = I_4,$$

$$\alpha_1 = \lambda_{1,2}, \quad \alpha_2 = \lambda_{2,4}, \quad \theta = 2\alpha_1 + 4\alpha_2. \tag{5.4}$$

With $Y_d Y_d^T = [z_{ij}^{(d)}]$, equation (3.10) becomes

$$\begin{bmatrix} z_{11}^{(1)} & z_{11}^{(1)} & z_{12}^{(1)} & z_{12}^{(1)} \\ z_{11}^{(1)} & z_{11}^{(1)} & z_{12}^{(1)} & z_{12}^{(1)} \\ z_{12}^{(1)} & z_{12}^{(1)} & z_{22}^{(1)} & z_{22}^{(1)} \\ z_{12}^{(1)} & z_{12}^{(1)} & z_{22}^{(1)} & z_{22}^{(1)} \end{bmatrix} + \begin{bmatrix} z_{11}^{(2)} + z_{22}^{(2)} & 0 & z_{13}^{(2)} + z_{24}^{(2)} & z_{14}^{(2)} + z_{23}^{(2)} \\ 0 & z_{11}^{(2)} + z_{22}^{(2)} & z_{14}^{(2)} + z_{23}^{(2)} & z_{13}^{(2)} + z_{24}^{(2)} \\ z_{13}^{(2)} + z_{24}^{(2)} & z_{14}^{(2)} + z_{23}^{(2)} & z_{33}^{(2)} + z_{44}^{(2)} & 0 \\ z_{14}^{(2)} + z_{23}^{(2)} & z_{13}^{(2)} + z_{24}^{(2)} & 0 & z_{33}^{(2)} + z_{44}^{(2)} \end{bmatrix} = 0,$$

i.e.,

$$\begin{cases} z_{11}^{(1)} = z_{22}^{(1)} = 0, \\ z_{11}^{(2)} + z_{22}^{(2)} = 0, \\ z_{33}^{(2)} + z_{44}^{(2)} = 0, \\ z_{12}^{(1)} + z_{14}^{(2)} + z_{23}^{(2)} = 0, \\ z_{12}^{(1)} + z_{13}^{(2)} + z_{24}^{(2)} = 0. \end{cases}$$

Thus, $L = 6$ and

$$C_1^{(1)} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \qquad C_1^{(2)} = 0,$$

$$C_2^{(1)} = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}, \qquad C_2^{(2)} = 0,$$

$$C_3^{(1)} = 0, \qquad C_3^{(2)} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix},$$

$$C_4^{(1)} = 0, \qquad C_4^{(2)} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix},$$

$$C_5^{(1)} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \qquad C_5^{(2)} = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix},$$

$$C_6^{(1)} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \qquad C_6^{(2)} = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

We have

$$\text{type}(a_1C_1^{(1)}+\cdots+a_6C_6^{(1)})=\begin{cases}(2,0,0,0) & \text{if } a_5=a_6,\ (a_1,a_2)=(0,0),\\ (2,0,0,1) & \text{if } a_5=a_6,\ (a_1,a_2)\ne(0,0),\\ (2,1,0,0) & \text{if } a_5\ne a_6,\ (a_1,a_2)\ne(1,1),\\ (2,1,1,0) & \text{if } a_5\ne a_6,\ (a_1,a_2)=(1,1),\end{cases}$$

$$\text{type}(a_1C_1^{(2)}+\cdots+a_6C_6^{(2)})=\begin{cases}(4,0,0,0) & \text{if } (a_5,a_6)=(0,0),\ (a_3,a_4)=(0,0),\\ (4,0,0,1) & \text{if } (a_5,a_6)=(0,0),\ (a_3,a_4)\ne(0,0),\\ (4,2,0,0) & \text{if } a_5\ne a_6,\\ (4,1,0,0) & \text{if } (a_5,a_6)=(1,1),\ (a_3,a_4)\ne(1,1),\\ (4,1,1,0) & \text{if } (a_5,a_6)=(1,1),\ (a_3,a_4)=(1,1).\end{cases}$$

As $(a_1,\dots,a_6)$ runs over $\mathbb{F}_2^6$,

$$\text{type}(\mathfrak{C}_{(a_1,\dots,a_6)})=\begin{cases}(\theta,0,0,0) & 1\text{ time},\\ (\theta,0,0,\delta(\alpha_2)) & 3\text{ times},\\ (\theta,\alpha_2,0,0) & 3\text{ times},\\ (\theta,\alpha_2,\alpha_2,0) & 1\text{ time},\\ (\theta,0,0,\delta(\alpha_1)) & 3\text{ times},\\ (\theta,0,0,\delta(\alpha_1+\alpha_2)) & 3^2\text{ times},\\ (\theta,\alpha_2,0,\delta(\alpha_1)) & 3^2\text{ times},\\ (\theta,\alpha_2,\alpha_2,\delta(\alpha_1)) & 3\text{ times},\\ (\theta,\alpha_1+2\alpha_2,0,0) & 2^3\cdot 3\text{ times},\\ (\theta,\alpha_1+2\alpha_2,\alpha_1,0) & 2^3\text{ times}.\end{cases}$$

By (3.18),

$$\begin{aligned}\mathfrak{n}(A_7)=\,&2^{2\alpha_1+4\alpha_2-6}\big[16-3\delta(\alpha_2)-3\delta(\alpha_1)-9\delta(\alpha_1+\alpha_2)\big]\\ &+2^{2\alpha_1+3\alpha_2-6}\big(3+(-1)^{\alpha_2}\big)\big(4-3\delta(\alpha_1)\big)+2^{\alpha_1+2\alpha_2-3}\big(3+(-1)^{\alpha_1}\big).\end{aligned}$$

By (3.9) and (5.4),

$$\begin{aligned}&|\text{Fix}(A_7,P_\lambda)|\\ &=2^{4\sum_{\nu(i)>1}\lambda_i}\,\mathfrak{n}(A_7)\\ &=2^{4\lambda_{0,4}}\Big[2^{2\lambda_{1,2}+4\lambda_{2,4}-6}\big[16-3\delta(\lambda_{2,4})-3\delta(\lambda_{1,2})-9\delta(\lambda_{1,2}+\lambda_{2,4})\big]\\ &\quad+2^{2\lambda_{1,2}+3\lambda_{2,4}-6}\big(3+(-1)^{\lambda_{2,4}}\big)\big(4-3\delta(\lambda_{1,2})\big)+2^{\lambda_{1,2}+2\lambda_{2,4}-3}\big(3+(-1)^{\lambda_{1,2}}\big)\Big].\end{aligned}$$

5.7. **Computation of** $|\text{Fix}(A_8,P_\lambda)|$.

The result follows immediately from Corollary 3.10 and equations (3.20) and (3.22). We have

$$\begin{aligned}|\text{Fix}(A_8,P_\lambda)|&=|\text{Fix}(\left[\begin{smallmatrix}0&1\\1&0\end{smallmatrix}\right],P_\lambda)|\,|\text{Fix}(\left[\begin{smallmatrix}0&1\\1&1\end{smallmatrix}\right],P_\lambda)|\\ &=2^{\lambda_{1,2}+2\lambda_{0,2}+2\lambda_{0,6}-3}\\ &\qquad\cdot\big[4-\delta(\lambda_{1,2})-\delta(\lambda_{2,4})-\delta(\lambda_{1,2}+\lambda_{2,4})\big]\big[2^{2\lambda_{3,6}}+(-1)^{\lambda_{3,6}}2^{\lambda_{3,6}}\big].\end{aligned}$$

5.8. **Computation of** $|\mathrm{Fix}(A_9, P_\lambda)|$.

We have $o(A_9) = 3$, $s_1 = 0$, $s_3 = 4$, $B_1 = \emptyset$, $B_3 = I_4$ and

$$\alpha_3 = \lambda_{3,6}. \tag{5.5}$$

With $Y_d Y_d^T = [z_{ij}^{(d)}]$, equation (3.10) becomes

$$\begin{bmatrix} 0 & z_{11}^{(3)} + z_{12}^{(3)} + z_{22}^{(3)} & z_{14}^{(3)} + z_{23}^{(3)} & z_{13}^{(3)} + z_{14}^{(3)} + z_{24}^{(3)} \\ z_{11}^{(3)} + z_{12}^{(3)} + z_{22}^{(3)} & 0 & z_{13}^{(3)} + z_{23}^{(3)} + z_{24}^{(3)} & z_{14}^{(3)} + z_{23}^{(3)} \\ z_{14}^{(3)} + z_{23}^{(3)} & z_{13}^{(3)} + z_{23}^{(3)} + z_{24}^{(3)} & 0 & z_{33}^{(3)} + z_{34}^{(3)} + z_{44}^{(3)} \\ z_{13}^{(3)} + z_{14}^{(3)} + z_{24}^{(3)} & z_{14}^{(3)} + z_{23}^{(3)} & z_{33}^{(3)} + z_{34}^{(3)} + z_{44}^{(3)} & 0 \end{bmatrix} = 0,$$

i.e.,

$$\begin{cases} z_{11}^{(3)} + z_{12}^{(3)} + z_{22}^{(3)} = 0, \\ z_{33}^{(3)} + z_{34}^{(3)} + z_{44}^{(3)} = 0, \\ z_{14}^{(3)} + z_{23}^{(3)} = 0, \\ z_{13}^{(3)} + z_{14}^{(3)} + z_{24}^{(3)} = 0. \end{cases}$$

Thus, $L = 4$ and

$$C_1^{(3)} = \begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad C_2^{(3)} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{bmatrix},$$

$$C_3^{(3)} = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad C_4^{(3)} = \begin{bmatrix} 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

As $(a_1, \dots, a_4)$ runs over $\mathbb{F}_2^4$,

$$\mathrm{type}(a_1 C_1^{(3)} + \cdots + a_4 C_4^{(3)}) = \begin{cases} (4,0,0,0) & 1 \text{ time}, \\ (4,1,1,0) & 5 \text{ times}, \\ (4,2,0,0) & 10 \text{ times}, \end{cases}$$

$$\mathrm{type}(\mathfrak{C}_{(a_1,\dots,a_4)}) = \begin{cases} (4\alpha_3,0,0,0) & 1 \text{ time}, \\ (4\alpha_3,\alpha_3,\alpha_3,0) & 5 \text{ times}, \\ (4\alpha_3,2\alpha_3,0,0) & 10 \text{ times}. \end{cases}$$

By (3.18),

$$\mathfrak{n}(A_9) = 2^{4\alpha_3 - 4} + 2^{3\alpha_3 - 4}\, 5(-1)^{\alpha_3} + 2^{2\alpha_3 - 3}\, 5.$$

By (3.9) and (5.5),

$$\begin{aligned} |\mathrm{Fix}(A_9, P_\lambda)| &= 2^{4\sum_{i \equiv 0\,(6)} \lambda_i}\, \mathfrak{n}(A_9) \\ &= 2^{4\lambda_{0,6} - 3} \big[2^{4\lambda_{3,6} - 1} + 2^{3\lambda_{3,6} - 1}\, 5(-1)^{\lambda_{3,6}} + 2^{2\lambda_{3,6}}\, 5 \big]. \end{aligned}$$

5.9. **Computation of $|\text{Fix}(A_{10}, P_\lambda)|$.**

We have $o(A_{10}) = 4$, $s_1 = 1$, $s_2 = 2$, $s_4 = 4$,

$$B_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \quad B_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad B_4 = I_4,$$

$$\alpha_1 = \lambda_{1,2}, \quad \alpha_2 = \lambda_{2,4}, \quad \alpha_4 = \lambda_{4,8}, \quad \theta = \alpha_1 + 2\alpha_2 + 4\alpha_4. \tag{5.6}$$

With $Y_d T_d^T = [z_{ij}^{(d)}]$, equation (3.10) becomes

$$\begin{bmatrix} z_{11}^{(1)} & z_{11}^{(1)} & z_{11}^{(1)} & z_{11}^{(1)} \\ z_{11}^{(1)} & z_{11}^{(1)} & z_{11}^{(1)} & z_{11}^{(1)} \\ z_{11}^{(1)} & z_{11}^{(1)} & z_{11}^{(1)} & z_{11}^{(1)} \\ z_{11}^{(1)} & z_{11}^{(1)} & z_{11}^{(1)} & z_{11}^{(1)} \end{bmatrix} + \begin{bmatrix} z_{11}^{(2)} + z_{22}^{(2)} & 0 & z_{11}^{(2)} + z_{22}^{(2)} & 0 \\ 0 & z_{11}^{(2)} + z_{22}^{(2)} & 0 & z_{11}^{(2)} + z_{22}^{(2)} \\ z_{11}^{(2)} + z_{22}^{(2)} & 0 & z_{11}^{(2)} + z_{22}^{(2)} & 0 \\ 0 & z_{11}^{(2)} + z_{22}^{(2)} & 0 & z_{11}^{(2)} + z_{22}^{(2)} \end{bmatrix}$$

$$+ \left[\begin{matrix} z_{11}^{(4)} + z_{22}^{(4)} + z_{33}^{(4)} + z_{44}^{(4)} & z_{12}^{(4)} + z_{23}^{(4)} + z_{34}^{(4)} + z_{14}^{(4)} \\ z_{12}^{(4)} + z_{23}^{(4)} + z_{34}^{(4)} + z_{14}^{(4)} & z_{11}^{(4)} + z_{22}^{(4)} + z_{33}^{(4)} + z_{44}^{(4)} \\ 0 & z_{12}^{(4)} + z_{23}^{(4)} + z_{34}^{(4)} + z_{14}^{(4)} \\ z_{12}^{(4)} + z_{23}^{(4)} + z_{34}^{(4)} + z_{14}^{(4)} & 0 \end{matrix}\right.$$

$$\left.\begin{matrix} 0 & z_{12}^{(4)} + z_{23}^{(4)} + z_{34}^{(4)} + z_{14}^{(4)} \\ z_{12}^{(4)} + z_{23}^{(4)} + z_{34}^{(4)} + z_{14}^{(4)} & 0 \\ z_{11}^{(4)} + z_{22}^{(4)} + z_{33}^{(4)} + z_{44}^{(4)} & z_{12}^{(4)} + z_{23}^{(4)} + z_{34}^{(4)} + z_{14}^{(4)} \\ z_{12}^{(4)} + z_{23}^{(4)} + z_{34}^{(4)} + z_{14}^{(4)} & z_{11}^{(4)} + z_{22}^{(4)} + z_{33}^{(4)} + z_{44}^{(4)} \end{matrix}\right] = 0,$$

i.e.,

$$\begin{cases} z_{11}^{(1)} + z_{11}^{(2)} + z_{22}^{(2)} = 0, \\ z_{11}^{(4)} + z_{22}^{(4)} + z_{33}^{(4)} + z_{44}^{(4)} = 0, \\ z_{11}^{(1)} + z_{12}^{(4)} + z_{23}^{(4)} + z_{34}^{(4)} + z_{14}^{(4)} = 0. \end{cases}$$

Thus, $L = 3$ and

$$C_1^{(1)} = [1], \qquad C_1^{(2)} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \qquad C_1^{(4)} = 0,$$

$$C_2^{(1)} = 0, \qquad C_2^{(2)} = 0, \qquad C_2^{(4)} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix},$$

$$C_3^{(1)} = [1], \qquad C_3^{(2)} = 0, \qquad C_2^{(4)} = \begin{bmatrix} 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

We have

$$\text{type}(a_1 C_1^{(1)} + a_2 C_2^{(1)} + a_3 C_3^{(1)}) = \begin{cases} (1,0,0,0) & \text{if } a_1 = a_3, \\ (1,0,0,1) & \text{if } a_1 \neq a_3, \end{cases}$$

$$\operatorname{type}(a_1C_1^{(2)}+a_2C_2^{(2)}+a_3C_3^{(2)})=\begin{cases}(2,0,0,0) & \text{if } a_1=0,\\ (2,0,0,1) & \text{if } a_1=1,\end{cases}$$

$$\operatorname{type}(a_1C_1^{(4)}+a_2C_2^{(4)}+a_3C_3^{(4)})=\begin{cases}(4,0,0,0) & \text{if } (a_2,a_3)=(0,0),\\ (4,1,0,0) & \text{if } (a_2,a_3)=(0,1),\\ (4,0,0,1) & \text{if } (a_2,a_3)=(1,0),\\ (4,1,1,0) & \text{if } (a_2,a_3)=(1,1).\end{cases}$$

As (a_1,a_2,a_3) runs over $\mathbb{F}_2^3$,

$$\operatorname{type}(\mathfrak{C}_{(a_1,a_2,a_3)})=\begin{cases}(\theta,0,0,0) & 1\text{ time},\\ (\theta,0,0,\delta(\alpha_4)) & 1\text{ time},\\ (\theta,\alpha_4,0,\delta(\alpha_2)) & 1\text{ time},\\ (\theta,\alpha_4,\alpha_4,\delta(\alpha_2)) & 1\text{ time},\\ (\theta,\alpha_4,0,\delta(\alpha_1)) & 1\text{ time},\\ (\theta,\alpha_4,\alpha_4,\delta(\alpha_1)) & 1\text{ time},\\ (\theta,0,0,\delta(\alpha_1+\alpha_2)) & 1\text{ time},\\ (\theta,0,0,\delta(\alpha_1+\alpha_2+\alpha_4)) & 1\text{ time}.\end{cases}$$

By (3.18),

$$\begin{aligned}\mathfrak{n}(A_{10})=&\,2^{\alpha_1+2\alpha_2+4\alpha_4-3}\bigl[4-\delta(\alpha_4)-\delta(\alpha_1+\alpha_2)-\delta(\alpha_1+\alpha_2+\alpha_4)\bigr]\\ &+2^{\alpha_1+2\alpha_2+3\alpha_4-3}\bigl(1+(-1)^{\alpha_4}\bigr)\bigl(2-\delta(\alpha_1)-\delta(\alpha_2)\bigr).\end{aligned}$$

By (3.9) and (5.6),

$$\begin{aligned}&|\operatorname{Fix}(A_{10},P_\lambda)|\\ &=2^{4\sum_{\nu(i)>2}\lambda_i}\,\mathfrak{n}(A_{10})\\ &=2^{4\lambda_{0,8}}\Bigl[2^{\lambda_{1,2}+2\lambda_{2,4}+4\lambda_{4,8}-3}\bigl[4-\delta(\lambda_{4,8})-\delta(\lambda_{1,2}+\lambda_{2,4})-\delta(\lambda_{1,2}+\lambda_{2,4}+\lambda_{4,8})\bigr]\\ &\quad+2^{\lambda_{1,2}+2\lambda_{2,4}+3\lambda_{4,8}-3}\bigl(1+(-1)^{\lambda_{4,8}}\bigr)\bigl(2-\delta(\lambda_{1,2})-\delta(\lambda_{2,4})\bigr)\Bigr].\end{aligned}$$

5.10. **Computation of** $|\operatorname{Fix}(A_{11},P_\lambda)|$**.**

We have $o(A_{11})=6$, $s_1=s_2=0$, $s_3=2$, $s_6=4$,

$$B_1=B_2=\emptyset,\qquad B_3=\begin{bmatrix}1&0\\1&1\\0&1\\1&0\end{bmatrix},\qquad B_6=I_4$$

and

$$\alpha_3=\lambda_{3,6},\qquad \alpha_6=\lambda_{6,12}\qquad \theta=2\alpha_3+4\alpha_6. \tag{5.7}$$

With $Y_dY_d^T=[z_{ij}^{(d)}]$, equation (3.10) becomes

$$\begin{bmatrix} 0 & z_{11}^{(3)}+z_{12}^{(3)}+z_{22}^{(3)} & z_{11}^{(3)}+z_{12}^{(3)}+z_{22}^{(3)} & 0 \\ z_{11}^{(3)}+z_{12}^{(3)}+z_{22}^{(3)} & 0 & z_{11}^{(3)}+z_{12}^{(3)}+z_{22}^{(3)} & z_{11}^{(3)}+z_{12}^{(3)}+z_{22}^{(3)} \\ z_{11}^{(3)}+z_{12}^{(3)}+z_{22}^{(3)} & z_{11}^{(3)}+z_{12}^{(3)}+z_{22}^{(3)} & 0 & z_{11}^{(3)}+z_{12}^{(3)}+z_{22}^{(3)} \\ 0 & z_{11}^{(3)}+z_{12}^{(3)}+z_{22}^{(3)} & z_{11}^{(3)}+z_{12}^{(3)}+z_{22}^{(3)}0 & \end{bmatrix}+$$

$$\begin{bmatrix} 0 & z_{12}^{(6)}+z_{14}^{(6)}+z_{34}^{(6)} & z_{11}^{(6)}+z_{13}^{(6)}+z_{22}^{(6)}+z_{24}^{(6)}+z_{33}^{(6)}+z_{44}^{(6)} & 0 \\ z_{12}^{(6)}+z_{14}^{(6)}+z_{34}^{(6)} & 0 & z_{12}^{(6)}+z_{14}^{(6)}+z_{34}^{(6)} & z_{11}^{(6)}+z_{13}^{(6)}+z_{22}^{(6)}+z_{24}^{(6)}+z_{33}^{(6)}+z_{44}^{(6)} \\ z_{11}^{(6)}+z_{13}^{(6)}+z_{22}^{(6)}+z_{24}^{(6)}+z_{33}^{(6)}+z_{44}^{(6)} & z_{12}^{(6)}+z_{14}^{(6)}+z_{34}^{(6)} & 0 & z_{12}^{(6)}+z_{14}^{(6)}+z_{34}^{(6)} \\ 0 & z_{11}^{(6)}+z_{13}^{(6)}+z_{22}^{(6)}+z_{24}^{(6)}+z_{33}^{(6)}+z_{44}^{(6)} & z_{12}^{(6)}+z_{14}^{(6)}+z_{34}^{(6)} & 0 \end{bmatrix}=0,$$

i.e.,

$$\begin{cases} z_{11}^{(3)}+z_{12}^{(3)}+z_{22}^{(3)}+z_{12}^{(6)}+z_{14}^{(6)}+z_{34}^{(6)}=0, \\ z_{11}^{(3)}+z_{12}^{(3)}+z_{22}^{(3)}+z_{11}^{(6)}+z_{13}^{(6)}+z_{22}^{(6)}+z_{24}^{(6)}+z_{33}^{(6)}+z_{44}^{(6)}=0. \end{cases}$$

So, $L=2$ and

$$C_1^{(3)}=\begin{bmatrix}1&1\\0&1\end{bmatrix}, \qquad C_1^{(6)}=\begin{bmatrix}0&1&0&1\\0&0&0&0\\0&0&0&1\\0&0&0&0\end{bmatrix},$$

$$C_2^{(3)}=\begin{bmatrix}1&1\\0&1\end{bmatrix}, \qquad C_2^{(6)}=\begin{bmatrix}1&0&1&0\\0&1&0&1\\0&0&1&0\\0&0&0&1\end{bmatrix}.$$

We have

$$\text{type}(a_1C_1^{(3)}+a_2C_2^{(3)})=\begin{cases}(1,0,0,0) & \text{if } a_1=a_2,\\ (1,1,1,0) & \text{if } a_1\ne a_2,\end{cases}$$

$$\text{type}(a_1C_1^{(6)}+a_2C_2^{(6)})=\begin{cases}(4,0,0,0) & \text{if } (a_1,a_2)=(0,0),\\ (4,2,0,0) & \text{if } a_1\ne a_2,\\ (4,1,1,0) & \text{if } (a_1,a_2)=(1,1),\end{cases}$$

$$\text{type}(\mathfrak{C}_{(a_1,a_2)})=\begin{cases}(\theta,0,0,0) & \text{if } (a_1,a_2)=(0,0),\\ (\theta,\alpha_3+2\alpha_6,\alpha_3,0) & \text{if } a_1\ne a_2,\\ (\theta,\alpha_6,\alpha_6,0) & \text{if } (a_1,a_2)=(1,1).\end{cases}$$

By (3.18),

$$\mathfrak{n}(A_{11})=2^{2\alpha_3+4\alpha_6-2}+2^{2\alpha_3+3\alpha_6-2}(-1)^{\alpha_6}+2^{\alpha_3+2\alpha_6-1}(-1)^{\alpha_3}.$$

By (3.9) and (5.7),

$$\begin{aligned}\mathrm{Fix}(A_{11}, P_\lambda) &= 2^{4\sum_{i\equiv 0\,(12)}\lambda_i}\,\mathfrak{n}(A_{11})\\ &= 2^{4\lambda_{0,12}-1}\big[2^{2\lambda_{3,6}+4\lambda_{6,12}-1}\\ &\quad + 2^{2\lambda_{3,6}+3\lambda_{6,12}-1}(-1)^{\lambda_{6,12}} + 2^{\lambda_{3,6}+2\lambda_{6,12}}(-1)^{\lambda_{3,6}}\big].\end{aligned}$$

5.11. **Computation of** $|\mathrm{Fix}(A_{12}, P_\lambda)|$**.**

We have $o(A_{12}) = 5$, $s_1 = 0$, $s_5 = 4$, $B_1 = \emptyset$, $B_5 = I_4$ and

$$\alpha_5 = \lambda_{5,10}. \tag{5.8}$$

With $Y_d Y_d^T = [z_{ij}^{(d)}]$, equation (3.10) becomes

$$\begin{bmatrix} 0 & z_{11}^{(5)} + z_{13}^{(5)} + z_{23}^{(5)} + z_{24}^{(5)} + z_{44}^{(5)} & z_{12}^{(5)} + z_{14}^{(5)} + z_{22}^{(5)} + z_{33}^{(5)} + z_{34}^{(5)} & z_{12}^{(5)} + z_{14}^{(5)} + z_{22}^{(5)} + z_{33}^{(5)} + z_{34}^{(5)}\\ z_{11}^{(5)} + z_{13}^{(5)} + z_{23}^{(5)} + z_{24}^{(5)} + z_{44}^{(5)} & 0 & z_{11}^{(5)} + z_{13}^{(5)} + z_{23}^{(5)} + z_{24}^{(5)} + z_{44}^{(5)} & z_{12}^{(5)} + z_{14}^{(5)} + z_{22}^{(5)} + z_{33}^{(5)} + z_{34}^{(5)}\\ z_{12}^{(5)} + z_{14}^{(5)} + z_{22}^{(5)} + z_{33}^{(5)} + z_{34}^{(5)} & z_{11}^{(5)} + z_{13}^{(5)} + z_{23}^{(5)} + z_{24}^{(5)} + z_{44}^{(5)} & 0 & z_{11}^{(5)} + z_{13}^{(5)} + z_{23}^{(5)} + z_{24}^{(5)} + z_{44}^{(5)}\\ z_{12}^{(5)} + z_{14}^{(5)} + z_{22}^{(5)} + z_{33}^{(5)} + z_{34}^{(5)} & z_{12}^{(5)} + z_{14}^{(5)} + z_{22}^{(5)} + z_{33}^{(5)} + z_{34}^{(5)} & z_{11}^{(5)} + z_{13}^{(5)} + z_{23}^{(5)} + z_{24}^{(5)} + z_{44}^{(5)} & 0 \end{bmatrix} = 0,$$

i.e.,

$$\begin{cases} z_{11}^{(5)} + z_{13}^{(5)} + z_{23}^{(5)} + z_{24}^{(5)} + z_{44}^{(5)} = 0,\\ z_{12}^{(5)} + z_{14}^{(5)} + z_{22}^{(5)} + z_{33}^{(5)} + z_{34}^{(5)} = 0. \end{cases}$$

Thus, $L = 2$ and

$$C_1^{(5)} = \begin{bmatrix} 1 & 0 & 1 & 0\\ 0 & 0 & 1 & 1\\ 0 & 0 & 0 & 0\\ 0 & 0 & 0 & 1 \end{bmatrix}, \qquad C_2^{(5)} = \begin{bmatrix} 0 & 1 & 0 & 1\\ 0 & 1 & 0 & 0\\ 0 & 0 & 1 & 1\\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

We have

$$\mathrm{type}(a_1 C_1^{(5)} + a_2 C_2^{(5)}) = \begin{cases} (4,0,0,0) & \text{if } (a_1,a_2) = (0,0),\\ (4,2,1,0) & \text{if } (a_1,a_2) \neq (0,0), \end{cases}$$

$$\mathrm{type}(\mathfrak{C}_{(a_1,a_2)}) = \begin{cases} (4\alpha_5,0,0,0) & \text{if } (a_1,a_2) = (0,0),\\ (4\alpha_5,2\alpha_5,\alpha_5,0) & \text{if } (a_1,a_2) \neq (0,0). \end{cases}$$

By (3.18), (3.9) and (5.8),

$$\mathfrak{n}(A_{12}) = 2^{4\alpha_5-2} + 2^{2\alpha_5-2}\,3(-1)^{\alpha_5},$$

$$\begin{aligned}|\mathrm{Fix}(A_{12}, P_\lambda)| &= 2^{4\sum_{i\equiv 0\,(10)}\lambda_i}\,\mathfrak{n}(A_{12})\\ &= 2^{4\lambda_{0,10}-2}\big[2^{4\lambda_{5,10}} + 2^{2\lambda_{5,10}}\,3(-1)^{\lambda_{5,10}}\big].\end{aligned}$$

5.12. Computation of $|\mathrm{Fix}(A_{13},P_\lambda)|$ and $|\mathrm{Fix}(A_{14},P_\lambda)|$.

The smallest positive integer t such that $x^4+x+1 \mid x^t-1$ is 15. It follows immediately from Lemma 3.11 that

$$|\mathrm{Fix}(A_{13},P_\lambda)| = 2^{4\lambda_{0,15}}. \tag{5.9}$$

In the same way,

$$|\mathrm{Fix}(A_{14},P_\lambda)| = 2^{4\lambda_{0,15}}. \tag{5.10}$$

6. Computation of $\Psi_{\le 5,n}$

The group $\mathrm{GL}(5,\mathbb{F}_2)$ has 27 conjugacy classes. A set of representatives, $A_1,\dots,A_{27}$, of the conjugacy classes is given in Table 3. In this section, we give explicit formulas for $|\mathrm{Fix}(A_i,P_\lambda)|$, $1\le i\le 27$. These formulas are computed by the same method in Sections 4 and 5. In many cases, Lemma 3.9 and the results of Sections 4 and 5 are used. The details of the computations are omitted.

$$\begin{aligned}|\mathrm{Fix}(A_1,P_\lambda)| = 2^{5\lambda_{0,2}}\Big[&2^{5\lambda_{1,2}-15}\big(18260-18259\,\delta(\lambda_{1,2})\big)\\ &+2^{4\lambda_{1,2}-15}\,155\big(3+(-1)^{\lambda_{1,2}}\big)+2^{3\lambda_{1,2}-12}\,217\big(5+3(-1)^{\lambda_{1,2}}\big)\Big].\end{aligned}$$

$$\begin{aligned}|\mathrm{Fix}(A_2,P_\lambda)| = 2^{5\lambda_{0,4}}\Big[&2^{4\lambda_{1,2}+5\lambda_{2,4}-11}\big[200-99\delta(\lambda_{1,2})-99\delta(\lambda_{1,2}+\lambda_{2,4})-\delta(\lambda_{2,4})\big]\\ &+2^{3\lambda_{1,2}+5\lambda_{2,4}-11}\,7\big(3+(-1)^{\lambda_{1,2}}\big)\big(2-\delta(\lambda_{2,4})\big)\\ &+2^{4\lambda_{1,2}+4\lambda_{2,4}-6}\,21\big(1-\delta(\lambda_{1,2})\big)+2^{3\lambda_{1,2}+4\lambda_{2,4}-8}\,7\big(3+(-1)^{\lambda_{1,2}}\big)\\ &+2^{2\lambda_{1,2}+4\lambda_{2,4}-7}\,7\big(5+3(-1)^{\lambda_{1,2}}\big)\Big].\end{aligned}$$

$$\begin{aligned}|\mathrm{Fix}(A_3,P_\lambda)| = 2^{5\lambda_{0,8}}\Big[&2^{3\lambda_{1,2}+4\lambda_{2,4}+5\lambda_{4,8}-7}\big[24-3\delta(\lambda_{1,2})-8\delta(\lambda_{1,2}+\lambda_{2,4})\\ &-3\delta(\lambda_{1,2}+\lambda_{2,4}+\lambda_{4,8})-8\delta(\lambda_{1,2}+\lambda_{4,8})-\delta(\lambda_{2,4}+\lambda_{4,8})\big]\\ &+2^{3\lambda_{1,2}+3\lambda_{2,4}+5\lambda_{4,8}-4}\,3\big[2-\delta(\lambda_{1,2})-\delta(\lambda_{1,2}+\lambda_{4,8})\big]\\ &+2^{2\lambda_{1,2}+4\lambda_{2,4}+5\lambda_{4,8}-7}\big(3+(-1)^{\lambda_{1,2}}\big)\big(2-\delta(\lambda_{2,4}+\lambda_{4,8})\big)\\ &+2^{2\lambda_{1,2}+3\lambda_{2,4}+5\lambda_{4,8}-6}\,3\big(3+(-1)^{\lambda_{1,2}}\big)\big(2-\delta(\lambda_{4,8})\big)\Big].\end{aligned}$$

$$\begin{aligned}|\mathrm{Fix}(A_4,P_\lambda)| = 2^{3\lambda_{0,2}+2\lambda_{0,6}-1}&\big(2^{2\lambda_{3,6}}+(-1)^{\lambda_{3,6}}2^{\lambda_{3,6}}\big)\\ &\cdot\big[2^{3\lambda_{1,2}-6}\big(36-35\,\delta(\lambda_{1,2})\big)+2^{2\lambda_{1,2}-6}\,7\big(3+(-1)^{\lambda_{1,2}}\big)\big].\end{aligned}$$

For $j=5,6$,

$$|\mathrm{Fix}(A_j,P_\lambda)| = 2^{2\lambda_{0,2}+3\lambda_{0,7}}\big[2^{2\lambda_{1,2}-3}\big(4-3\,\delta(\lambda_{1,2})\big)+2^{\lambda_{1,2}-3}\big(3+(-1)^{\lambda_{1,2}}\big)\big].$$

Table 3. Information about $\mathcal{C}(\mathrm{GL}(5,\mathbb{F}_2))$

representative	elementary divisors	$\lvert \mathrm{cent}_{\mathrm{GL}(5,\mathbb{F}_2)}(\ ) \rvert$
$A_1 = I_5$	$x+1,\ x+1,\ x+1,\ x+1,\ x+1$	$2^{10}\cdot 3^2\cdot 5\cdot 7\cdot 31$
$A_2 = I_3 \oplus \begin{bmatrix}0&1\\1&0\end{bmatrix}$	$x+1,\ x+1,\ x+1,\ (x+1)^2$	$2^{10}\cdot 3\cdot 7$
$A_3 = I_2 \oplus \begin{bmatrix}0&1&0\\0&0&1\\1&1&1\end{bmatrix}$	$x+1,\ x+1,\ (x+1)^3$	$2^7\cdot 3$
$A_4 = I_3 \oplus \begin{bmatrix}0&1\\1&1\end{bmatrix}$	$x+1,\ x+1,\ x+1,\ x^2+x+1$	$2^3\cdot 3^2\cdot 7$
$A_5 = I_2 \oplus \begin{bmatrix}0&1&0\\0&0&1\\1&1&0\end{bmatrix}$	$x+1,\ x+1,\ x^3+x+1$	$2\cdot 3\cdot 7$
$A_6 = I_2 \oplus \begin{bmatrix}0&1&0\\0&0&1\\1&0&1\end{bmatrix}$	$x+1,\ x+1,\ x^3+x^2+1$	$2\cdot 3\cdot 7$
$A_7 = [1] \oplus \begin{bmatrix}0&1\\1&0\end{bmatrix} \oplus \begin{bmatrix}0&1\\1&0\end{bmatrix}$	$x+1,\ (x+1)^2,\ (x+1)^2$	$2^9\cdot 3$
$A_8 = [1] \oplus \begin{bmatrix}0&1\\1&0\end{bmatrix} \oplus \begin{bmatrix}0&1\\1&1\end{bmatrix}$	$x+1,\ (x+1)^2,\ x^2+x+1$	$2^3\cdot 3$
$A_9 = [1] \oplus \begin{bmatrix}0&1\\1&1\end{bmatrix} \oplus \begin{bmatrix}0&1\\1&1\end{bmatrix}$	$x+1,\ x^2+x+1,\ x^2+x+1$	$2^2\cdot 3^2\cdot 5$
$A_{10} = [1] \oplus \begin{bmatrix}0&1&0&0\\0&0&1&0\\0&0&0&1\\1&0&0&0\end{bmatrix}$	$x+1,\ (x+1)^4$	2^5
$A_{11} = [1] \oplus \begin{bmatrix}0&1&0&0\\0&0&1&0\\0&0&0&1\\1&0&1&0\end{bmatrix}$	$x+1,\ (x^2+x+1)^2$	$2^2\cdot 3$
$A_{12} = [1] \oplus \begin{bmatrix}0&1&0&0\\0&0&1&0\\0&0&0&1\\1&1&1&1\end{bmatrix}$	$x+1,\ x^4+x^3+x^2+x+1$	$3\cdot 5$
$A_{13} = [1] \oplus \begin{bmatrix}0&1&0&0\\0&0&1&0\\0&0&0&1\\1&1&0&0\end{bmatrix}$	$x+1,\ x^4+x+1$	$3\cdot 5$
$A_{14} = [1] \oplus \begin{bmatrix}0&1&0&0\\0&0&1&0\\0&0&0&1\\1&0&0&1\end{bmatrix}$	$x+1,\ x^4+x^3+1$	$3\cdot 5$
$A_{15} = \begin{bmatrix}0&1\\1&0\end{bmatrix} \oplus \begin{bmatrix}0&1&0\\0&0&1\\1&1&1\end{bmatrix}$	$(x+1)^2,\ (x+1)^3$	2^7
$A_{16} = \begin{bmatrix}0&1\\1&0\end{bmatrix} \oplus \begin{bmatrix}0&1&0\\0&0&1\\1&1&0\end{bmatrix}$	$(x+1)^2,\ x^3+x+1$	$2\cdot 7$

Table 3. Continued

$A_{17} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \oplus \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 1 \end{bmatrix}$	$(x+1)^2,\ x^3+x^2+1$	$2 \cdot 7$
$A_{18} = \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix} \oplus \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 1 & 1 \end{bmatrix}$	$x^2+x+1,\ (x+1)^3$	$2^2 \cdot 3$
$A_{19} = \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix} \oplus \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix}$	$x^2+x+1,\ x^3+x+1$	$3 \cdot 7$
$A_{20} = \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix} \oplus \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 1 \end{bmatrix}$	$x^2+x+1,\ x^3+x^2+1$	$3 \cdot 7$
$A_{21} = \begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 1 & 1 & 0 & 0 & 1 \end{bmatrix}$	$(x+1)^5$	2^4
$A_{22} = \begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 & 0 \end{bmatrix}$	x^5+x^2+1	31
$A_{23} = \begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 & 0 \end{bmatrix}$	x^5+x^3+1	31
$A_{24} = \begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 1 & 1 & 1 & 1 & 0 \end{bmatrix}$	$x^5+x^3+x^2+x+1$	31
$A_{25} = \begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 1 & 1 & 1 & 0 & 1 \end{bmatrix}$	$x^5+x^4+x^2+x+1$	31
$A_{26} = \begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 1 & 1 & 0 & 1 & 1 \end{bmatrix}$	$x^5+x^4+x^3+x+1$	31
$A_{27} = \begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & 1 & 1 \end{bmatrix}$	$x^5+x^4+x^3+x^2+1$	31

$$\begin{aligned}|\mathrm{Fix}(A_7, P_\lambda)| = 2^{5\lambda_{0,4}} \Big[& 2^{3\lambda_{1,2}+5\lambda_{2,4}-9} \big[80 - 7\delta(\lambda_{1,2}) - 69\delta(\lambda_{1,2}+\lambda_{2,4}) - 3\delta(\lambda_{2,4})\big] \\ & + 2^{3\lambda_{1,2}+4\lambda_{2,4}-9} \big[72 + 8(-1)^{\lambda_{2,4}} - 7(-1)^{\lambda_{2,4}}\delta(\lambda_{1,2}) - 69\delta(\lambda_{1,2})\big] \\ & + 2^{2\lambda_{1,2}+5\lambda_{2,4}-7}\, 3\big(3+(-1)^{\lambda_{1,2}}\big)\big(1-\delta(\lambda_{2,4})\big) \\ & + 2^{2\lambda_{1,2}+4\lambda_{2,4}-7}\, 3\big(3+(-1)^{\lambda_{1,2}}\big) + 2^{3\lambda_{1,2}+3\lambda_{2,4}-2}\big(1-\delta(\lambda_{1,2})\big) \\ & + 2^{2\lambda_{1,2}+3\lambda_{2,4}-4}\big(3+(-1)^{\lambda_{1,2}}\big)\Big].\end{aligned}$$

$$\begin{aligned}|\mathrm{Fix}(A_8, P_\lambda)| = {} & 2^{3\lambda_{0,4}+2\lambda_{0,6}-1}\big(2^{2\lambda_{3,6}} + (-1)^{\lambda_{3,6}} 2^{\lambda_{3,6}}\big) \\ & \cdot \Big[2^{2\lambda_{1,2}+3\lambda_{2,4}-4}\big[8 - 3\delta(\lambda_{1,2}) - \delta(\lambda_{2,4}) - 3\delta(\lambda_{1,2}+\lambda_{2,4})\big] \\ & + 2^{\lambda_{1,2}+2\lambda_{2,4}-3}\big(3+(-1)^{\lambda_{1,2}}\big)\Big].\end{aligned}$$

$$\begin{aligned}|\mathrm{Fix}(A_9, P_\lambda)| = {} & 2^{\lambda_{0,1}+4\lambda_{0,6}-4}\big(2-\delta(\lambda_{1,2})\big) \\ & \cdot \big[2^{4\lambda_{3,6}-1} + 2^{3\lambda_{3,6}-1}\, 5(-1)^{\lambda_{3,6}} + 2^{2\lambda_{3,6}}\, 5\big].\end{aligned}$$

$$\begin{aligned}|\mathrm{Fix}(A_{10}, P_\lambda)| = 2^{5\lambda_{0,8}} \Big[& 2^{2\lambda_{1,2}+3\lambda_{2,4}+5\lambda_{4,8}-5}\big[8 - \delta(\lambda_{1,2}) - 2\delta(\lambda_{1,2}+\lambda_{2,4}) \\ & - 2\delta(\lambda_{1,2}+\lambda_{2,4}+\lambda_{4,8}) - \delta(\lambda_{1,2}+\lambda_{4,8}) - \delta(\lambda_{4,8})\big] \\ & + 2^{2\lambda_{1,2}+3\lambda_{2,4}+4\lambda_{4,8}-5}\big(1+(-1)^{\lambda_{4,8}}\big) \\ & \cdot \big[4 - 2\delta(\lambda_{1,2}) - \delta(\lambda_{1,2}+\lambda_{2,4}) - \delta(\lambda_{2,4})\big] \\ & + 2^{\lambda_{1,2}+2\lambda_{2,4}+5\lambda_{4,8}-4}\big(3+(-1)^{\lambda_{1,2}}\big)\big(1-\delta(\lambda_{4,8})\big) \\ & + 2^{\lambda_{1,2}+2\lambda_{2,4}+4\lambda_{4,8}-4}\big(3+(-1)^{\lambda_{1,2}}\big)\Big].\end{aligned}$$

$$\begin{aligned}|\mathrm{Fix}(A_{11}, P_\lambda)| = {} & 2^{\lambda_{0,1}+4\lambda_{0,12}-2}\big(2-\delta(\lambda_{1,2})\big)\big[2^{2\lambda_{3,6}+4\lambda_{6,12}-1} \\ & + 2^{2\lambda_{3,6}+3\lambda_{6,12}-1}(-1)^{\lambda_{6,12}} + 2^{\lambda_{3,6}+2\lambda_{6,12}}(-1)^{\lambda_{3,6}}\big].\end{aligned}$$

$$|\mathrm{Fix}(A_{12}, P_\lambda)| = 2^{\lambda_{0,1}+4\lambda_{0,10}-3}\big(2-\delta(\lambda_{1,2})\big)\big[2^{4\lambda_{5,10}} + 2^{2\lambda_{5,10}}\, 3(-1)^{\lambda_{5,10}}\big].$$

For $j = 13, 14$,

$$|\mathrm{Fix}(A_j, P_\lambda)| = 2^{\lambda_{0,1}+4\lambda_{0,15}-1}\big(2-\delta(\lambda_{1,2})\big).$$

$$\begin{aligned}|\mathrm{Fix}(A_{15},P_\lambda)| = 2^{5\lambda_{0,8}}\Big[&2^{2\lambda_{1,2}+4\lambda_{2,4}+5\lambda_{4,8}-6}\big[16-\delta(\lambda_{1,2})-5\delta(\lambda_{1,2}+\lambda_{2,4})\\ &-4\delta(\lambda_{1,2}+\lambda_{2,4}+\lambda_{4,8})-2\delta(\lambda_{1,2}+\lambda_{4,8})\\ &-\delta(\lambda_{2,4})-2\delta(\lambda_{2,4}+\lambda_{4,8})\big]\\ &+2^{2\lambda_{1,2}+3\lambda_{2,4}+5\lambda_{4,8}-6}\big[12+4(-1)^{\lambda_{2,4}}-2(-1)^{\lambda_{2,4}}\delta(\lambda_{1,2})\\ &-4\delta(\lambda_{1,2})-5\delta(\lambda_{1,2}+\lambda_{4,8})-(-1)^{\lambda_{2,4}}\delta(\lambda_{1,2}+\lambda_{4,8})\\ &-\delta(\lambda_{4,8})-(-1)^{\lambda_{2,4}}\delta(\lambda_{4,8})\big]\\ &+2^{\lambda_{1,2}+2\lambda_{2,4}+4\lambda_{4,8}-3}\big(3+(-1)^{\lambda_{1,2}}\big)\Big].\end{aligned}$$

For $j=16,17$,

$$|\mathrm{Fix}(A_j,P_\lambda)| = 2^{\lambda_{1,2}+2\lambda_{0,2}+3\lambda_{0,7}-2}\big[4-\delta(\lambda_{1,2})-\delta(\lambda_{2,4})-\delta(\lambda_{1,2}+\lambda_{2,4})\big].$$

$$\begin{aligned}|\mathrm{Fix}(A_{18},P_\lambda)| = 2^{2\lambda_{0,6}+\lambda_{1,2}+2\lambda_{2,4}+3\lambda_{0,4}-3}&\big[2^{2\lambda_{3,6}}+(-1)^{\lambda_{3,6}}2^{\lambda_{3,6}}\big]\\ &\cdot\big[4-\delta(\lambda_{1,2}+\lambda_{2,4})-\delta(\lambda_{1,2}+\lambda_{4,8})-\delta(\lambda_{2,4}+\lambda_{4,8})\big].\end{aligned}$$

For $j=19,20$,

$$|\mathrm{Fix}(A_j,P_\lambda)| = 2^{2\lambda_{0,6}+3\lambda_{0,7}-1}\big[2^{2\lambda_{3,6}}+(-1)^{\lambda_{3,6}}2^{\lambda_{3,6}}\big].$$

$$\begin{aligned}|\mathrm{Fix}(A_{21},P_\lambda)| = 2^{5\lambda_{0,16}}\Big[&2^{\lambda_{1,2}+2\lambda_{2,4}+4\lambda_{4,8}+5\lambda_{8,16}-3}\big[4-\delta(\lambda_{1,2}+\lambda_{2,4})\\ &-\delta(\lambda_{1,2}+\lambda_{2,4}+\lambda_{4,8})-\delta(\lambda_{4,8})\big]\\ &+2^{\lambda_{1,2}+2\lambda_{2,4}+3\lambda_{4,8}+5\lambda_{8,16}-3}\big(1+(-1)^{\lambda_{4,8}}\big)\\ &\cdot\big[2-\delta(\lambda_{1,2}+\lambda_{8,16})-\delta(\lambda_{2,4}+\lambda_{8,16})\big]\Big].\end{aligned}$$

For $22\le j\le 27$,

$$|\mathrm{Fix}(A_j,P_\lambda)| = 2^{5\lambda_{0,31}}.$$

7. NUMERICAL RESULTS

Two tables are included in this section. Table 4 contains the values of $\Psi_{\le k,n}$ with $k\le 5$ and $n\le 40$. Table 5 contains the values of $\Psi_{k,n}$ with k and n in the same ranges

REFERENCES

[1] J. H. Conway and V. Pless, *On the enumeration of self-dual codes*, J. Combin. Theory A **28** (1980), 26 – 53.

[2] J. H. Conway, V. Pless, and N. J. A. Sloane, *The binary self-dual codes of length up to 32: a revised enumeration*, J. Combin. Theory A **60** (1992), 183 – 195.

[3] L. E. Dickson, *Linear Groups*, Dover, New York, 1958.

Table 4. Values of $\Psi_{\le k,n}$, $k \le 5$, $n \le 40$

$n\backslash k$	0	1	2	3	4	5
1	1	1	1	1	1	1
2	1	2	2	2	2	2
3	1	2	2	2	2	2
4	1	3	4	4	4	4
5	1	3	4	4	4	4
6	1	4	7	8	8	8
7	1	4	7	9	9	9
8	1	5	11	16	18	18
9	1	5	11	17	20	20
10	1	6	16	28	37	39
11	1	6	16	30	42	46
12	1	7	23	49	77	92
13	1	7	23	53	89	112
14	1	8	31	82	157	218
15	1	8	31	89	187	281
16	1	9	41	133	323	551
17	1	9	41	144	389	740
18	1	10	53	210	654	1,447
19	1	10	53	229	804	2,059
20	1	11	67	325	1,324	4,029
21	1	11	67	354	1,651	6,032
22	1	12	83	490	2,654	11,774
23	1	12	83	534	3,356	18,581
24	1	13	102	727	5,291	36,239
25	1	13	102	793	6,759	59,798
26	1	14	123	1,058	10,433	116,020
27	1	14	123	1,154	13,444	198,489
28	1	15	147	1,515	20,363	382,272
29	1	15	147	1,651	26,384	670,031
30	1	16	174	2,136	39,229	1,276,454
31	1	16	174	2,329	51,025	2,267,431
32	1	17	204	2,972	74,574	4,260,828
33	1	17	204	3,237	97,143	7,596,889
34	1	18	237	4,078	139,660	14,050,410
35	1	18	237	4,439	181,923	24,965,555
36	1	19	274	5,532	257,592	45,384,782
37	1	19	274	6,017	335,029	79,965,507
38	1	20	314	7,418	467,600	142,792,476
39	1	20	314	8,061	606,613	248,697,834
40	1	21	358	9,843	835,392	497,412,483

TABLE 5. Values of $\Psi_{k,n}$, $k \le 5$, $n \le 40$

$n\backslash k$	0	1	2	3	4	5
1	1	0	0	0	0	0
2	1	1	0	0	0	0
3	1	1	0	0	0	0
4	1	2	1	0	0	0
5	1	2	1	0	0	0
6	1	3	3	1	0	0
7	1	3	3	2	0	0
8	1	4	6	5	2	0
9	1	4	6	6	3	0
10	1	5	10	12	9	2
11	1	5	10	14	12	4
12	1	6	16	26	28	15
13	1	6	16	30	36	23
14	1	7	23	51	75	61
15	1	7	23	58	98	94
16	1	8	32	92	190	228
17	1	8	32	103	245	351
18	1	9	43	157	444	793
19	1	9	43	176	575	1,255
20	1	10	56	258	999	2,705
21	1	10	56	287	1,297	4,381
22	1	11	71	407	2,164	9,120
23	1	11	71	451	2,822	15,225
24	1	12	89	625	4,564	30,948
25	1	12	89	691	5,966	53,039
26	1	13	109	935	9,375	105,587
27	1	13	109	1,031	12,290	185,045
28	1	14	132	1,368	18,848	361,909
29	1	14	132	1,504	24,733	643,647
30	1	15	158	1,962	37,093	1,237,225
31	1	15	158	2,155	48,696	2,216,406
32	1	16	187	2,768	71,602	4,186,254
33	1	16	187	3,033	93,906	7,499,746
34	1	17	219	3,841	135,582	13,910,750
35	1	17	219	4,202	177,484	24,783,632
36	1	18	255	5,258	252,060	45,127,190
37	1	18	255	5,743	329,012	79,630,478
38	1	19	294	7,104	460,182	142,324,876
39	1	19	294	7,747	598,552	248,091,221
40	1	20	337	9,485	825,549	496,577,091

[4] J. F. Dillon and H. Dobbertin, *New cyclic difference set with Singer parameters*, Finite Fields Appl. **10** (2004), 342 – 389.

[5] X. Hou, *Classification of cosets of the Reed-Muller code* $R(m-3,m)$, Discrete Math. **128** (1994), 203 – 224.

[6] X. Hou, $\mathrm{GL}(m,2)$ *acting on* $R(r,m)/R(r-1,m)$, Discrete Math. **149** (1996), 99 – 122.

[7] X. Hou, *Affinity of permutations of* $\mathbb{F}_2^n$, Discrete Appl. Math. **154** (2006), 313 – 325.

[8] X. Hou, *On the asymptotic number of inequivalent binary self-dual codes*, J. Combin. Theory A, Available online 22 August 2006.

[9] F. J. MacWilliams and N. J. A. Sloane *The Theory of Error-Correcting Codes*, North-Holland, New York, 1981

[10] V. Pless, *A classification of self-orthogonal codes over* GF(2), Discrete Math. **3** (1972), 209 – 246.

[11] V. Pless and N. J. A. Sloane, *On the classification and enumeration of self-dual codes*, J. Combin. Theory A **18** (1975), 313 – 335.

Department of Mathematics, University of South Florida, Tampa, FL 33620
E-mail address: `xhou@tarski.math.usf.edu`